

Nondegenerate cohomology pairing for transitive Lie algebroids, characterization

Jan Kubarski^{1*}, Alexandr Mishchenko^{2†}

¹ *Institute of Mathematics,
Technical University of Łódź,
al. Politechniki 11, 90-924 Łódź, Poland*

² *Department of Mathematics,
Moscow State University,
Leninskije Gory 119992, Moscow, Russia*

Received 15 December 2003; accepted 15 April 2004

Abstract: The Evens-Lu-Weinstein representation (Q_A, D) for a Lie algebroid A on a manifold M is studied in the transitive case. To consider at the same time non-oriented manifolds as well, this representation is slightly modified to (Q_A^{or}, D^{or}) by tensoring by orientation flat line bundle, $Q_A^{or} = Q_A \otimes or(M)$ and $D^{or} = D \otimes \partial_A^{or}$. It is shown that the induced cohomology pairing is nondegenerate and that the representation (Q_A^{or}, D^{or}) is the unique (up to isomorphism) line representation for which the top group of compactly supported cohomology is nontrivial. In the case of trivial Lie algebroid $A = TM$ the theorem reduce to the following: the orientation flat bundle $(or(M), \partial^{or})$ is the unique (up to isomorphism) flat line bundle (ξ, ∇) for which the twisted de Rham complex of compactly supported differential forms on M with values in ξ possesses the nontrivial cohomology group in the top dimension. Finally it is obtained the characterization of transitive Lie algebroids for which the Lie algebroid cohomology with trivial coefficients (or with coefficients in the orientation flat line bundle) gives Poincaré duality. In proofs of these theorems for Lie algebroids it is used the Hochschild-Serre spectral sequence and it is shown the general fact concerning pairings between graded filtered differential \mathbb{R} -vector spaces: assuming that the second terms live in the finite rectangular, nondegeneration of the pairing for the second terms (which can be infinite dimensional) implies the same for cohomology spaces.

© Central European Science Journals. All rights reserved.

Keywords: twisted cohomology, cohomology of Lie algebras, Poincaré duality, Lie algebroid, modular class, cohomology pairing, Evens-Lu-Weinstein pairing, Hochschild-Serre spectral sequence

MSC (2000): 58 H 99, 17 B 56, 18 G 40, 55 R 20, 55 T 05, 57 R 19, 57 R 22, 58 A 10

* E-mail: kubarski@mail.p.lodz.pl

† E-mail: asmish@higeom.math.msu.su

1 Introduction

The cohomology pairing coming from Evens-Lu-Weinstein representation of a Lie algebroid [4] is very important in many applications of Lie algebroids (Poisson geometry, intrinsic characteristic classes). This pairing generalizes the well known pairings that give Poincaré duality for Lie algebra cohomology and de Rham cohomology of a manifold and real cohomology of transitive invariantly oriented Lie algebroids [14]. For a Poisson manifold, this pairing agree with the pairing on the Poisson homology. The authors of [4] give an example of a nontransitive Lie algebroid for which the pairing is not necessarily nondegenerate and post the problem of when it is nondegenerate. This paper gives the positive answer for the case of any transitive Lie algebroids and proves the property of this representation: it is the one (up to isomorphism) for which the top group of compactly supported cohomology is nontrivial.

Finally, we prove that for the nonregular transformation Lie algebroid corresponding to the action $\gamma : \mathbb{R} \rightarrow \mathfrak{X}(\mathbb{R})$, $\gamma(t) = t \cdot X$ where $X = x^N \frac{d}{dx}$, there is no line representation for which the cohomological pairing is nondegenerate.

More detailed, this paper is devoted to prove two cycles of theorems, mutually overcoming.

FIRST CYCLE concerns nondegenerate cohomology pairings for manifolds (Theorem 2.3), Lie algebras (Theorem 3.4) and Lie algebroids (Theorem 7.3).

— Assume that M is a connected m -dimensional manifold (oriented or not) and ξ_1, ξ_2 are two flat vector bundles with flat covariant derivatives ∇_1 and ∇_2 respectively. Denote by $or(M)$ the orientation bundle with canonical flat structure ∂^{or} . Let $F : \xi_1 \times \xi_2 \rightarrow or(M)$ be a pairing (i.e. 2-linear homomorphism) of vector bundles compatible with the flat structures $(\nabla_1, \nabla_2, \partial^{or})$, nondegenerate at least at one point (therefore, at every). From such a pairing one obtains a pairing on differential forms and the induced pairing in cohomology

$$H_{\nabla_1}^j(M, \xi_1) \times H_{\nabla_2, c}^{m-j}(M, \xi_2) \xrightarrow{F\#} H_{\partial^{or}, c}^m(M, or(M)) \xrightarrow{J_M^{or, \#}} \mathbb{R}$$

is nondegenerate in the sense that

$$H_{\nabla_1}^j(M, \xi_1) \xrightarrow{\cong} (H_{\nabla_2, c}^{m-j}(M, \xi_2))^*.$$

The index "c" means that the compactly supported cohomology are considered. This theorem generalizes the classical Poincaré duality as well as the one for d_ω -cohomologies [6].

— Assume that \mathfrak{g} is an arbitrary n -dimensional Lie algebra and $\nabla_1, \nabla_2 : \mathfrak{g} \rightarrow L_{\mathbb{R}}$ are two representations of \mathfrak{g} in \mathbb{R} . Denote by $\nabla_{\text{trad}} : \mathfrak{g} \rightarrow L_{\mathbb{R}}$ the trace-representation $(\nabla_{\text{trad}})_a = \text{tr}(\text{ad}_a) \cdot \text{id}$. Then the top group of cohomology $H_{\text{trad}}^n(\mathfrak{g})$ of \mathfrak{g} with respect to ∇_{trad} is nonzero, $H_{\text{trad}}^n(\mathfrak{g}) \xrightarrow{\cong} \mathbb{R}$, and if the multiplication of reals is compatible with respect to $(\nabla_1, \nabla_2, \nabla_{\text{trad}})$ then the exterior multiplication $\wedge : \Lambda^r \mathfrak{g}^* \times \Lambda^{n-r} \mathfrak{g}^* \rightarrow \Lambda^n \mathfrak{g}^* \cong \mathbb{R}$ yields the induced nondegenerate pairing in cohomology

$$H_{\nabla_1}^i(\mathfrak{g}) \times H_{\nabla_2}^{n-i}(\mathfrak{g}) \rightarrow H_{\text{trad}}^n(\mathfrak{g}) \cong \mathbb{R},$$

i.e.

$$H^i_{\nabla_1}(\mathfrak{g}) \xrightarrow{\cong} (H^{n-i}_{\nabla_2}(\mathfrak{g}))^* .$$

In particular, for $(\nabla_1, \nabla_2, \nabla_{\text{trad}}) = (0, \nabla_{\text{trad}}, \nabla_{\text{trad}})$ we obtain

$$H^i(\mathfrak{g}) \xrightarrow{\cong} (H^{n-i}_{\text{trad}}(\mathfrak{g}))^* .$$

For unimodular Lie algebra \mathfrak{g} the usual Poincaré duality is obtained in this way.

— Let $A = (A, [\cdot, \cdot], \#_A)$ be a Lie algebroid on M and

$$Q_A = \Lambda^{\text{top}} A \otimes \Lambda^{\text{top}} T^* M$$

the line vector bundle with canonical Evens-Lu-Weinstein representation [4],

$$D_\gamma(Y \otimes \varphi) = L_\gamma(Y) \otimes \varphi + Y \otimes L_{\#_A(\gamma)}(\varphi) .$$

To consider non-oriented manifolds we modify it into

$$Q_A^{or} = Q_A \otimes or(M)$$

and

$$D^{or} = D \otimes \partial_A^{or}$$

tensoring by the orientation bundle and its flat structure $((\partial_A^{or})_\gamma \sigma = (\partial^{or})_{\#_A(\gamma)} \sigma, \sigma \in \Gamma(or(M)), \#_A : A \rightarrow TM$ is the anchor of A). For transitive Lie algebroids with n -dimensional isotropy Lie algebras and multiplications by reals $(M \times \mathbb{R}) \otimes Q_A^{or} \rightarrow Q_A^{or}$ the induced pairing in cohomology

$$H^j(A) \times H^{m+n-j}_{D^{or,c}}(A, Q_A^{or}) \rightarrow H^{m+n}_{D^{or,c}}(A, Q_A^{or}) \rightarrow \mathbb{R}$$

is nondegenerate, i.e. $H^{m+n}_{D^{or,c}}(A, Q_A^{or}) \cong \mathbb{R}$ and

$$H^j(A) \cong (H^{m+n-j}_{D^{or,c}}(A, Q_A^{or}))^* .$$

SECOND CYCLE shows the uniqueness of the line representation for which the top group of compactly supported cohomology is not zero (Theorems 2.10, 3.5, 7.10).

— $H^m_{\nabla,c}(M, \xi) \neq 0$ if and only if (ξ, ∇) is, up to isomorphism, the orientation flat line bundle $(or(M), \partial^{or})$. In particular, for oriented manifold, $H^m_{\nabla,c}(M, \xi) \neq 0$ if and only if (ξ, ∇) is, up to isomorphism, the trivial flat line bundle $(M \times \mathbb{R}, \partial)$.

— For an n -dimensional Lie algebra \mathfrak{g} the trace-representation ∇_{trad} is the unique line representation ∇ for which $H^n_{\nabla}(\mathfrak{g}) \neq 0$.

— Let A be a transitive Lie algebroid and ∇ a representation of A in a line vector bundle ξ . Then $H^{m+n}_{\nabla,c}(A, \xi) \neq 0$ if and only if (ξ, ∇) is, up to isomorphism, the E-L-W-representation (Q_A^{or}, D^{or}) .

In conclusion we obtain a full classification of transitive Lie algebroids for which the algebra of real cohomologies with trivial coefficients satisfies the Poincaré duality.

— If A is a transitive Lie algebroid then the following conditions are equivalent:

- $H_c^{m+n}(A) \neq 0$,
- $H_c^{m+n}(A) \cong \mathbb{R}$ and $H^j(A) \cong (H_c^{m+n-j}(A))^*$,
- A is orientable vector bundle and the modular class of A is zero, $\theta_A = 0$.

In particular,

— For an orientable manifold M we have: $H_c^{m+n}(A) \neq 0$ if and only if A is a TUIO-Lie algebroid [13], i.e. the adjoint Lie Algebra Bundle $\mathfrak{g} = \ker \#_A$ is oriented and there is a global nonsingular section $\varepsilon \in \Gamma(\Lambda^n \mathfrak{g})$ invariant with respect to the adjoint representation.

The above theorem for a compact oriented manifold M and 1-rank adjoint LAB $\mathfrak{g} = M \times \mathbb{R}$ was proved earlier in [15].

To prove Theorem 7.10 we use Theorem 4.4 concerning a pairing $\cdot : {}^1A \times {}^2A \rightarrow {}^3A$ between graded filtered differential \mathbb{R} -vector spaces and their spectral sequences. Roughly speaking, if the second terms ${}^rE_2^{j,i}$ live in the rectangular $0 \leq j \leq m, 0 \leq i \leq n$, ${}^3E_2^{(m+n)} = {}^3E_2^{m,n} \cong \mathbb{R}$ and the multiplication of the second terms $\langle \cdot, \cdot \rangle_2 : {}^1E_2^{(j)} \times {}^2E_2^{(m+n-j)} \rightarrow {}^3E_2^{m,n} \cong \mathbb{R}$ is nondegenerate in the sense that ${}^1E_2^{(j)} \cong ({}^2E_2^{(m+n-j)})^*$, then the cohomology pairing for cohomologies of spaces is nondegenerate as well, i.e. ${}^3H^{m+n} \cong \mathbb{R}$ and ${}^1H^j \xrightarrow{\cong} ({}^2H^{m+n-j})^*$. We must stress that the spaces ${}^rE_2^{j,i}$ may be infinite dimensional.

2 Non-degenerate pairings for twisted cohomology of a manifold

Many of the facts from this section belong to "the folklore". We call 1-dimensional vector bundles *line bundles*.

2.1 Twisted cohomology, elementary properties

Let M be an m -dimensional paracompact manifold and ξ a vector bundle of rank p and $\nabla_X \nu$, $X \in \mathfrak{X}(M)$, $\nu \in \Gamma(\xi)$, a **flat** covariant derivative on M in ξ .

- (•) The differential equation $\nabla \nu = 0$ (with respect to the local section ν of ξ) is locally uniquely integrable.

The local section ν satisfying $\nabla \nu = 0$ is called ∇ -*constant* (or sometimes ∇ -*invariant*). To set a flat covariant derivative ∇ is equivalent to set local trivializations $\{(U_\alpha, \varphi_\alpha)\}$ relative to which the transitive functions are locally constant which is, in turn, equivalent to set a homomorphism of Lie algebroids $\nabla : TM \rightarrow A(\xi)$ where $A(\xi)$ is the Lie algebroid of ξ . The flat bundles (ξ, ∇) form a category with morphisms $F : (\xi_1, \nabla_1) \rightarrow (\xi_2, \nabla_2)$ being linear isomorphisms $F : \xi_1 \rightarrow \xi_2$ compatible with flat covariant derivatives (∇_1, ∇_2) , i.e. for which $F(\nabla_{1,X}\nu) = \nabla_{2,X}(F\nu)$. We write also

$$(\xi_1, \nabla_1) \stackrel{F}{\sim} (\xi_2, \nabla_2)$$

or briefly $\nabla_1 \stackrel{F}{\sim} \nabla_2$.

Two flat line bundles over a connected manifold are isomorphic if and only if they have the same holonomy homomorphism $h : \pi_1(M, x) \rightarrow GL(\mathbb{R}, 1)$. For a flat vector bundle (ξ, ∇) the differential operator d_∇ of the degree 1 for ξ -valued differential forms $\Omega^*(M, \xi)$ is defined by the standard formula

$$d_\nabla(\phi)(X_0, \dots, X_q) = \sum_i (-1)^i \nabla_{X_i}(\phi(X_0, \dots, \hat{i}, \dots, X_q)) + \sum_{i < j} (-1)^{i+j} \phi([X_i, X_j], X_0, \dots, \hat{i}, \dots, \hat{j}, \dots, X_q).$$

Let $\sigma_\alpha^1, \dots, \sigma_\alpha^p$ be local sections of ξ over U_α corresponding to the standard basis $e^1, \dots, e^p \in \mathbb{R}^p$ under the trivialization φ_α , $\sigma_\alpha^i(x) = \varphi_{\alpha,x}(e^i)$. The local sections σ_α^i are ∇ -constant, $\nabla\sigma_\alpha^i = 0$. Over U_α a ξ -valued q -form ϕ can be written as $\sum \phi^i \otimes \sigma_\alpha^i$, $\phi^i \in \Omega^q(U_\alpha)$ and we have $d_\nabla(\sum \phi^i \otimes \sigma_\alpha^i) = \sum d_{dR}(\phi^i) \otimes \sigma_\alpha^i$. The flatness of ∇ implies that d_∇ is a differential operator, $d_\nabla^2 = 0$, therefore $\Omega^*(M, \xi)$ is a differential complex and the (twisted) cohomology

$$H_\nabla^*(M, \xi) = H(\Omega^*(M, \xi), d_\nabla)$$

makes sense. By the definition the 0-group of cohomology can be written as

$$H_\nabla^0(M, \xi) = \{\nu \in \Gamma(\xi); \nabla_X \nu = 0 \text{ for all } X \in \mathfrak{X}(M)\}. \tag{1}$$

(••) If (ξ, ∇) is a line nontrivial flat vector bundle then according to (•) above $H_\nabla^0(M, \xi) = 0$.

If ∇ is a flat covariant derivative in a vector bundle ξ and $\omega \in \Omega^1(M)$ is a closed real 1-form, then

$$\nabla_X^\omega \nu = \nabla_X \nu + \omega(X) \cdot \nu \tag{2}$$

is a flat covariant derivative as well. If ξ is a line bundle and ∇ and ∇_1 are two flat covariant derivatives then there exists a closed 1-form ω such that $\nabla_1 = \nabla^\omega$.

Each flat covariant derivative ∇ in the trivial vector bundle $M \times \mathbb{R}$ is of the form ∂^ω for some closed 1-form ω (∂ is the standard covariant derivative in the trivial vector bundle $M \times \mathbb{R}$ defined by differentiation of functions $\partial_X(f) = X(f)$). Differential operator d_{∂^ω} is given directly by

$$d_{\partial^\omega}(\phi) = d_{dR}\phi + \omega \wedge \phi.$$

The operator d_{∂^ω} is in the literature denoted rather by d_ω than by d_{∂^ω} [6], [8] and the cohomology space $H_{\partial^\omega}(M, M \times \mathbb{R})$ is denoted by $H_\omega(M)$. If $\omega = 0$ the usual de Rham cohomology of M is obtained. It is easy to see that

$$H_\omega^0(M) = \begin{cases} 0 & \iff \omega \text{ is nonexact,} \\ \mathbb{R} & \iff \omega \text{ is exact.} \end{cases} \tag{3}$$

The space of ξ -valued q -forms with compact support $\Omega_c^*(M, \xi)$ is a differential complex as well and we have the compactly supported cohomology $H_{\nabla,c}^*(M, \xi)$. If $(\xi_1, \nabla_1) \overset{F}{\sim} (\xi_2, \nabla_2)$ then $F_* : \Omega^*(M, \xi_1) \rightarrow \Omega^*(M, \xi_2)$ commutes with the differential operators d_{∇_1}

and d_{∇_2} and gives rise to an isomorphism in cohomology $F_{\#} : H_{\nabla_1}^*(M, \xi_1) \rightarrow H_{\nabla_2}^*(M, \xi_2)$. Analogously, for compact supports, we have an isomorphism $F_{\#,c} : H_{\nabla_1,c}^*(M, \xi_1) \rightarrow H_{\nabla_2,c}^*(M, \xi_2)$.

For an open subset $U \subset M$ we have the restricted flat covariant derivative ∇_U on U in the vector bundle ξ_U and the twisted cohomology $H_{\nabla}(U, \xi)$ and $H_{\nabla,c}(U, \xi)$ are defined. Similarly as in the case of real coefficients (see for example [1]) we can obtain the short exact Mayer-Vietoris sequences ($U_1, U_2 \subset M$ are open subset, $U = U_1 \cup U_2$, and $U_{12} = U_1 \cap U_2$)

$$0 \rightarrow \Omega^*(U, \xi) \xrightarrow{\alpha} \Omega^*(U_1, \xi) \oplus \Omega^*(U_2, \xi) \xrightarrow{\beta} \Omega^*(U_{12}, \xi) \rightarrow 0$$

and

$$0 \leftarrow \Omega_c^*(U, \xi) \xleftarrow{\alpha_c} \Omega_c^*(U_1, \xi) \oplus \Omega_c^*(U_2, \xi) \xleftarrow{\beta_c} \Omega_c^*(U_{12}, \xi) \leftarrow 0.$$

They give rise to long exact sequences in cohomology

$$\rightarrow H_{\nabla}^q(U, \xi) \xrightarrow{\alpha_{\#}} H_{\nabla}^q(U_1, \xi) \oplus H_{\nabla}^q(U_2, \xi) \xrightarrow{\beta_{\#}} H_{\nabla}^q(U_{12}, \xi) \xrightarrow{\partial^q} H_{\nabla}^{q+1}(U, \xi) \rightarrow$$

and

$$\leftarrow H_{\nabla,c}^q(U, \xi) \xleftarrow{\alpha_{c\#}} H_{\nabla,c}^q(U_1, \xi) \oplus H_{\nabla,c}^q(U_2, \xi) \xleftarrow{\beta_{c\#}} H_{\nabla,c}^q(U_{12}, \xi) \xleftarrow{\partial_c^q} H_{\nabla,c}^{q+1}(U, \xi) \leftarrow$$

Remark 2.1. There is a natural isomorphism $H_{\nabla}^*(M, \xi) \cong H_{I(\nabla)}^*(\mathfrak{U}, \xi)$ of $H_{\nabla}^*(M, \xi)$ with $H_{I(\nabla)}^*(\mathfrak{U}, \xi)$, the cohomology of M in the sheaf $I(\nabla)$ of local ∇ -constant sections of ξ . In other words, $H_{\nabla}^*(M, \xi)$ are cohomology of M with local system of coefficients.

2.2 Orientation flat bundle and its characterization

Let $\{(U_{\alpha}, x_{\alpha})\}$ be a coordinate open cover for the manifold M , with transition functions $g_{\alpha\beta} = x_{\alpha} \circ x_{\beta}^{-1}$. Take the orientation bundle $or(M)$, i.e. the line bundle on M with a distinguished system of local trivializations $\{\varphi_{\alpha}\}$ such that the transition functions are equal to $\text{sgn } J(g_{\alpha\beta})$ [1]. Let $\{e_{\alpha}\}$ be a family of local sections corresponding to 1 under the trivializations $\{\varphi_{\alpha}\}$, $e_{\alpha}(x) = \varphi_{\alpha,x}(1)$. In the bundle $or(M)$ there exists exactly one flat covariant derivative ∂^{or} such that e_{α} are ∂^{or} -constant, $\partial^{or}(e_{\alpha}) = 0$. The notation e_{α} and ∂^{or} is valid in the whole paper.

The flat orientation bundle $(or(M), \partial^{or})$ is characterized by the holonomy homomorphism $s : \pi_1(M, x_0) \rightarrow \mathbb{Z}_2 \subset GL(\mathbb{R}, 1)$ that can be identified with monodromy to the group of the local orientations in the fixed point x_0 which also is \mathbb{Z}_2 .

In the sequel it will be useful to give other characterization of the flat orientation bundle.

Proposition 2.2. *Let (ξ, ∇) be a flat line bundle. The following conditions are equivalent:*

(a) $(\xi, \nabla) \cong (or(M), \partial^{or}),$

- (b) there exists a collection of local sections $\{\sigma_\alpha\}$ of ξ such that σ_α are ∇ -constant and the transition functions are equal to $\text{sgn } J(g_{\alpha\beta})$,
- (c) $(or(M) \otimes \xi, \partial^{or} \otimes \nabla) \cong (M \times \mathbb{R}, \partial)$,
- (d) there exists a global nonsingular section $t \in \Gamma(or(M) \otimes \xi)$ which is $\partial^{or} \otimes \nabla$ -constant.

Proof. Equivalences (a) \Leftrightarrow (b) and (c) \Leftrightarrow (d) are evident by definition.

(b) \Rightarrow (c) The linear homomorphism $F : or(M) \otimes \xi \rightarrow M \times \mathbb{R}$ defined by $F(e_\alpha \otimes \sigma_\alpha) = 1$ is a well defined linear isomorphism compatible with $(\partial^{or} \otimes \nabla, \partial)$.

(d) \Rightarrow (b) Locally $t = e_\alpha \otimes \sigma_\alpha$ for some local nonsingular sections σ_α of ξ . Since

$$0 = \partial^{or} \otimes \nabla (e_\alpha \otimes \sigma_\alpha) = \partial^{or} \otimes \sigma_\alpha + e_\alpha \otimes \nabla \sigma_\alpha = e_\alpha \otimes \nabla \sigma_\alpha,$$

it follows that σ_α are ∇ -constant and have the same transition functions $\text{sgn } J(g_{\alpha\beta})$.

The $or(M)$ -valued m -differential forms are called *densities*. There exists an operator

$$\int_M^{or} : \Omega_c^m(M, or(M)) \rightarrow \mathbb{R}$$

of the integration of densities and the Stoke’s Theorem for densities holds

$$\int_M^{or} d_{\partial^{or}}(\omega) = 0$$

for $\omega \in \Omega_c^{m-1}(M, or(M))$ [1]. Hence it produces a linear operator

$$\int_M^{or, \#} : H_{\partial^{or}, c}^m(M, or(M)) \rightarrow \mathbb{R}. \tag{4}$$

2.3 Pairings and cohomology, nondegeneracy

Now let (ξ_1, ∇_1) , (ξ_2, ∇_2) , and (ξ_3, ∇_3) be three flat vector bundles. We say that (ξ_1, ∇_1) and (ξ_2, ∇_2) are *paired* to (ξ_3, ∇_3) if there is a bilinear homomorphism $F : \xi_1 \times \xi_2 \rightarrow \xi_3$ compatible with flat covariant derivatives $(\nabla_1, \nabla_2, \nabla_3)$, i.e. such that, for every $X \in \mathfrak{X}(M)$,

$$\nabla_{3, X} F(\nu_1, \nu_2) = F(\nabla_{1, X} \nu_1, \nu_2) + F(\nu_1, \nabla_{2, X} \nu_2) .. \tag{5}$$

Then we write $F : (\xi_1, \nabla_1) \times (\xi_2, \nabla_2) \rightarrow (\xi_3, \nabla_3)$. From such a pairing one obtains a pairing $(\phi, \psi) \mapsto \phi \wedge \psi := F_*(\phi, \psi)$ of $\Omega^q(M, \xi_1)$ and $\Omega^r(M, \xi_2)$ to $\Omega^{q+r}(M, \xi_3)$ fullfilling the equality

$$d_{\nabla_3} F_*(\phi, \psi) = F_*(d_{\nabla_1} \phi, \psi) + (-1)^{\text{deg } \phi} F_*(\phi, d_{\nabla_2} \psi).$$

Clearly, $\phi \wedge \psi := F_*(\phi, \psi)$ is the usual wedge product of differential forms with F -multiplication of values, see [7, Vol.II]. The pairing of differential forms induces a pairing of cohomology classes

$$F_{\#} : H_{\nabla_1}^*(M, \xi_1) \times H_{\nabla_2}^*(M, \xi_2) \rightarrow H_{\nabla_3}^*(M, \xi_3)$$

as well as the pairing for compact supports

$$F_{\#,c} : H_{\nabla_1}^*(M, \xi_1) \times H_{\nabla_2,c}^*(M, \xi_2) \rightarrow H_{\nabla_3,c}^*(M, \xi_3).$$

Consider two flat vector bundles (ξ_1, ∇_1) , (ξ_2, ∇_2) and a pairing

$$F : (\xi_1, \nabla_1) \times (\xi_2, \nabla_2) \rightarrow (or(M), \partial^{or}). \quad (6)$$

For an open subset $U \subset M$ we define a pairing

$$\int_U^{or,\#} \circ F_{\#} : H_{\nabla_1}^q(U, \xi_1) \times H_{\nabla_2,c}^{m-q}(U, \xi_2) \xrightarrow{F_{\#}} H_{\partial^{or},c}^m(U, or(M)) \xrightarrow{\int_U^{or,\#}} \mathbb{R},$$

and the Poincaré linear homomorphism

$$D_U^q : H_{\nabla_1}^q(U, \xi_1) \rightarrow (H_{\nabla_2,c}^{m-q}(U, \xi_2))^*, \quad D_U^q([\Phi])([\Psi]) = \int_U (\Phi \wedge \Psi).$$

Similarly as in the case of real coefficients we check that the family of Poincaré homomorphisms $\{D_U^q\}$ induces a map from the long exact sequences in cohomology to the long exact sequences in compactly supported cohomology (the symbols of vector bundles ξ_1 and ξ_2 in the diagram below are omitted and the sign \pm is equal precisely to $(-1)^{q+1}$)

$$\begin{array}{ccccccc} H_{\nabla}^q(U) & \xrightarrow{\alpha_{\#}} & H_{\nabla}^q(U_1) \oplus H_{\nabla}^q(U_2) & \xrightarrow{\beta_{\#}} & H_{\nabla}^q(U_{12}) & \xrightarrow{\partial^q} & H_{\nabla}^{q+1}(U) \\ \downarrow D_U & & \downarrow D_{U_1 \oplus U_2} & & \downarrow D_{U_{12}} & & \downarrow D_U \\ H_{\nabla,c}^q(U, \xi)^* & \xrightarrow{\alpha_{c\#}^*} & H_{\nabla,c}^q(U_1)^* \oplus H_{\nabla,c}^q(U_2)^* & \xrightarrow{\beta_{c\#}^*} & H_{\nabla,c}^q(U_{12})^* & \xrightarrow{\pm(\partial_c^q)^*} & H_{\nabla,c}^{q+1}(U)^* \end{array} \quad (7)$$

For an infinite disjoint open subsets $U = \coprod U_i$ we deduce that D_U can be identifying with $\prod D_{U_i}$.

Theorem 2.3. *Assume that M is connected. If pairing (6) is nondegenerate at least one point then the **cohomology pairing***

$$\int_M^{or,\#} \circ F_{\#} : H_{\nabla_1}^q(M, \xi_1) \times H_{\nabla_2,c}^{m-q}(M, \xi_2) \xrightarrow{F_{\#}} H_{\partial^{or},c}^m(M, or(M)) \xrightarrow{\int_M^{or,\#}} \mathbb{R},$$

is also nondegenerate in the sense that

$$D_M^q : H_{\nabla_1}^q(M, \xi_1) \xrightarrow{\cong} (H_{\nabla_2,c}^{m-q}(M, \xi_2))^*$$

is an isomorphism, $q \in \{0, 1, \dots, m\}$.

Proof. We can use the standard method from [7, Vol.I] (or a slightly modified method by using Riemannian structure and properties of geodesically convex neighbourhoods, [1], [18]). According to [7, Vol.I, Prop.II, p.16] and the commutativity of diagram (7) and remark on infinite disjoint open subsets we need only to prove the theorem for the manifold $M = \mathbb{R}^m$.

Each vector bundle ξ over \mathbb{R}^m is trivial, each flat covariant derivative ∇ has trivial holonomy, so the differential equation $\nabla\nu = 0$ is globally integrable. Therefore for an arbitrary point $x_0 \in M$ there exists an isomorphism of flat vector bundles

$$\varphi : (\xi, \nabla) \rightarrow (\mathbb{R}^m \times \xi_{x_0}, \partial)$$

where by ∂ is denoted the standard flat covariant derivative $\partial_X f = X(f)$.

(Remark: for the line bundle ξ the isomorphism φ can be given directly as follows. For $\xi = \mathbb{R}^m \times \mathbb{R}$ any flat covariant derivative ∇ is of the form $\nabla_X f = \partial_X f + \partial_X(\alpha) \cdot f$ for some function α . Then $\varphi(f) = e^{-\alpha} f$ is a required isomorphism.)

The isomorphism φ gives rise to an isomorphism in cohomology

$$\varphi_{\#} : H_{\nabla}(\mathbb{R}^m, \xi) \xrightarrow{\cong} H_{dR}(\mathbb{R}^m, \xi_{x_0}),$$

especially for the zero level

$$H_{\nabla}^0(\mathbb{R}^m, \xi) \xrightarrow[\cong]{\varphi_{\#}^0} H_{dR}^0(\mathbb{R}^m, \xi_{x_0}) \xrightarrow[\cong]{\rho} \xi_{x_0}.$$

On the other hand, the isomorphism φ also gives rise to an isomorphism in compactly supported cohomology $\varphi_{\#,c} : H_{\nabla,c}(\mathbb{R}^m, \xi) \xrightarrow{\cong} H_{dR,c}(\mathbb{R}^m, \xi_{x_0})$, especially for the top level

$$\varphi_{\#,c}^m : H_{\nabla,c}^m(\mathbb{R}^m, \xi) \xrightarrow[\cong]{\varphi_{\#,c}^m} H_{dR,c}^m(\mathbb{R}^m, \xi_{x_0}) \xrightarrow[\cong]{\rho_c} \xi_{x_0},$$

where ρ_c is defined by the formula

$$\rho_c([\sum_i f^i \cdot \Delta \otimes v_i]) = \sum_i (\int_{\mathbb{R}^m} f^i) \cdot e^i$$

where v_i is a basis of ξ_{x_0} , Δ is a determinant function on \mathbb{R}^m and $f^i \in C_c^\infty(\mathbb{R}^m)$ are functions with compact support. ρ_c is independent of the choice of the basis v_i and fulfils the equality $\rho_c([f \cdot \Delta \otimes v]) = (\int_{\mathbb{R}^m} f) \cdot v$, $f \in C_c^\infty(\mathbb{R}^m)$, $v \in \xi_{x_0}$.

Now take flat vector bundles (ξ_i, ∇_i) on \mathbb{R}^m and linear isomorphisms $\varphi_i : (\xi_i, \nabla_i) \rightarrow (\mathbb{R}^m \times (\xi_i)_{x_0}, \partial)$. For any pairing $F : (\xi_1, \nabla_1) \times (\xi_2, \nabla_2) \rightarrow (\xi_3, \nabla_3)$ we get easily the commutative diagram

$$\begin{array}{ccc} F_{\#} : H_{\nabla_1}^0(\mathbb{R}^m, \xi_1) \times H_{\nabla_2,c}^m(\mathbb{R}^m, \xi_2) & \longrightarrow & H_{\nabla_3,c}^m(\mathbb{R}^m, \xi_3) \\ \downarrow \varphi_{1\#}^0 \times \varphi_{2\#,c}^m & & \downarrow \varphi_{3\#,c}^m \\ \bar{F}_{\#} : H_{dR}^0(\mathbb{R}^m, (\xi_1)_{x_0}) \times H_{dR,c}^m(\mathbb{R}^m, (\xi_2)_{x_0}) & \longrightarrow & H_{dR,c}^m(\mathbb{R}^m, (\xi_3)_{x_0}) \\ \downarrow \rho \times \rho_c & & \downarrow \rho_c \\ F_{x_0} : (\xi_1)_{x_0} \times (\xi_2)_{x_0} & \longrightarrow & (\xi_3)_{x_0} \end{array}$$

where the middle pairing comes from the "constant" pairing

$$\bar{F} : (\mathbb{R}^m, (\xi_1)_{x_0}) \times (\mathbb{R}^m, (\xi_2)_{x_0}) \rightarrow (\mathbb{R}^m, (\xi_3)_{x_0}), \quad \bar{F}_x = F_{x_0}.$$

To prove the theorem take $(\xi_3, \nabla_3) = (or(M), \partial^{or})$ and choose a point x_0 such that F_{x_0} is nondegenerate.

2.4 Applications of the nondegenerate cohomology pairing

Now we give a number of applications of Theorem 2.3.

Example 2.4. For a connected orientable manifold M and the trivial flat vector bundles $(\xi_i, \nabla_i) = (M \times \mathbb{R}, \partial)$ and the multiplication of reals $\cdot : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ we obtain the classical Poincaré duality $H^j(M) \times H_c^{m-j}(M) \rightarrow H_c^m(M) \rightarrow \mathbb{R}$. Especially $H_c^m(M) = \mathbb{R}$ and $H^j(M) \cong (H_c^{m-j}(M))^*$.

Example 2.5. More generally, for arbitrary connected manifold M taking $(\xi_1, \nabla_1) = (M \times \mathbb{R}, \partial)$ and $(\xi_2, \nabla_2) = (or(M), \partial^{or})$ and the multiplication by reals $F : (M \times \mathbb{R}) \times or(M) \rightarrow or(M)$ we get the Poincaré duality also for nonorientable manifold [1]. Especially operator (4) is an isomorphism, $H_{\partial^{or},c}^m(M, or(M)) \cong \mathbb{R}$.

Example 2.6. [6], [8] Let M be an oriented connected manifold. The following conditions are equivalent:

- (1) $H_{\omega,c}^m(M) = 0$,
- (2) $H^1(M) \ni [\omega] \neq 0$.

If $[\omega] = 0$ then $H_{\omega,c}^m(M) = \mathbb{R}$.

Indeed, consider multiplication by reals $F : (M \times \mathbb{R}) \times (M \times \mathbb{R}) \rightarrow M \times \mathbb{R}$. This pairing is nondegenerate and compatible with $(\partial^{-\omega}, \partial^{\omega}, \partial)$. By Theorem 2.3 we get the nondegenerate pairing $H_{-\omega}^p(M) \times H_{\omega,c}^{m-p}(M) \rightarrow H_c^m(M) \cong \mathbb{R}$. In particular, we get $H_{-\omega}^0(M) = (H_{\omega,c}^m(M))^*$, so all follows from (3).

Each flat covariant derivative in $or(M)$ is of the form $(\partial^{or})^{\omega}$ for a closed 1-form ω . Consider the multiplications by reals $F : (M \times \mathbb{R}) \times or(M) \rightarrow or(M)$. Then we easily get:

Example 2.7. For any connected manifold M (oriented or not) the following conditions are equivalent:

- (1) $H_{(\partial^{or})^{\omega},c}^m(M, or(M)) = 0$,
- (2) $H^1(M) \ni [\omega] \neq 0$.

If $[\omega] = 0$ then $H_{(\partial^{or})^{\omega},c}^m(M, or(M)) = \mathbb{R}$.

The next applications are given in the following propositions.

Proposition 2.8. *If M is orientable and ξ is an arbitrary line nonorientable (i.e. non-trivial) vector bundle then for any flat covariant derivative ∇ in ξ*

$$H_{\nabla,c}^m(M, \xi) = 0.$$

Proof. Indeed, consider the natural nondegenerate pairing

$$F : (\xi, \nabla) \times (\xi, \nabla) \rightarrow (\xi \otimes \xi, \nabla \otimes \nabla), \quad (\nu, \mu) \mapsto \nu \otimes \mu,$$

and any linear isomorphism $\varphi : \xi \otimes \xi \rightarrow M \times \mathbb{R}$. The latter transforms the flat covariant derivative $\nabla \otimes \nabla$ to the ∂^ω for some closed 1-form ω . We recall that $(\nabla \otimes \nabla)_X (\nu \otimes \mu) = \nabla_X \nu \otimes \mu + \nu \otimes \nabla_X \mu$. Then the pairing $\varphi \circ F : \xi \times \xi \rightarrow M \times \mathbb{R}$ is compatible with $(\nabla, \nabla, \partial^\omega)$ and, in consequence, with $(\nabla^{-\omega}, \nabla, \partial)$ (for $\nabla^{-\omega}$ see (2)). By Theorem 2.3 we have the nondegenerate pairing

$$H_{\nabla^{-\omega}}^0(M, \xi) \times H_{\nabla, c}^m(M, \xi) \rightarrow H_c^m(M) \xrightarrow{f_M} \mathbb{R}.$$

In consequence we obtain by the nontriviality of ξ and observation (•) from section 2.1

$$0 = H_{\nabla^{-\omega}}^0(M, \xi) = (H_{\nabla, c}^m(M, \xi))^*$$

which imply $H_{\nabla, c}^m(M, \xi) = 0$.

Proposition 2.9. *If ξ is a line bundle not isomorphic to $or(M)$ then for arbitrary flat covariant derivative ∇ in ξ we have*

$$H_{\nabla, c}^m(M, \xi) = 0.$$

Proof. Indeed, fix a linear isomorphism

$$\varphi : \xi \otimes \xi \rightarrow M \times \mathbb{R}.$$

Such isomorphism φ exists since $\xi \otimes \xi$ is orientable line vector bundle, therefore, trivial. Let $\nabla \otimes \nabla \stackrel{F}{\sim} \partial^\omega$ for a closed 1-form ω . Take the multiplication by reals

$$\tau : or(M) \otimes (M \times \mathbb{R}) \rightarrow or(M)$$

and notice that τ is compatible with $(\partial^{or} \otimes \partial^\omega, (\partial^{or})^\omega)$. Consider the canonical nondegenerate pairing $F : (or(M) \otimes \xi) \times \xi \rightarrow or(M) \otimes \xi \otimes \xi$ which is compatible with $(\partial^{or} \otimes \nabla, \nabla, \partial^{or} \otimes \nabla \otimes \nabla)$. The composition

$$F' : (or(M) \otimes \xi) \times \xi \xrightarrow{F} or(M) \otimes \xi \otimes \xi \xrightarrow{id \otimes \varphi} or(M) \otimes (M \times \mathbb{R}) \xrightarrow{\tau} or(M),$$

clearly, is nondegenerate and is compatible with $(\partial^{or} \otimes \nabla, \nabla, (\partial^{or})^\omega)$. Therefore F' is compatible with $(\partial^{or} \otimes \nabla^{-\omega}, \nabla, \partial^{or})$. According to Theorem 2.3 applied to F' we get

$$H_{\partial^{or} \otimes \nabla^{-\omega}}^0(M, or(M) \otimes \xi) \cong (H_{\nabla, c}^m(M, \xi))^*.$$

Since ξ is not isomorphic to $or(M)$ the vector bundle $or(M) \otimes \xi$ is not trivial (indeed, if $or(M) \otimes \xi \cong M \times \mathbb{R}$ then $or(M) \cong \xi^* \cong \xi$) which produces $H_{\partial^{or} \otimes \nabla^{-\omega}}^0(M, or(M) \otimes \xi) = 0$ and further $H_{\nabla, c}^m(M, \xi) = 0$.

Finally we have the main application.

Theorem 2.10. *The following conditions are equivalent:*

- a) $H_{\nabla, c}^m(M, \xi) \neq 0$,
 b) $H_{\nabla, c}^m(M, \xi) = \mathbb{R}$,
 c) $(\xi, \nabla) \sim (or(M), \partial^{or})$,

Proof. For c) \implies b) see Example 2.5 or [1]; b) \implies a) is evident. It remains to show that

a) \implies c). Keep the notation φ and ω from the proof of the previous proposition. By the same reasoning we check

$$H_{\partial^{or} \otimes \nabla^{-\omega}}^0(M, or(M) \otimes \xi) \neq 0.$$

It means that $or(M) \otimes \xi$ is trivial and there exists a nonsingular global cross-section $\nu \in \Gamma(or(M) \otimes \xi)$ which is $\partial^{or} \otimes \nabla^{-\omega}$ -constant. Express locally ν in the form $\nu = e_\alpha \otimes f_\alpha$ for some local sections f_α of ξ , for e_α see subsection 2.2. It is evident that $\{f_\alpha\}$ has the transition function equal to $\text{sgn } Jg_{\alpha\beta}$ and that $\nabla^{-\omega} f_\alpha = 0$, i.e. $\nabla_X f_\alpha = \omega(X) \cdot f_\alpha$. The formula $f = \varphi(f_\alpha \otimes f_\alpha)$ determines correctly a nonsingular function f . Since $\nabla \otimes \nabla \stackrel{\mathcal{L}}{\sim} \partial^\omega$ then $\partial_X f + \omega(X) \cdot f = \partial_X^\omega f = 2 \cdot \omega(X) \cdot f$, one has

$$\partial_X f = \omega(X) \cdot f.$$

The global cross-section $\nu' = \frac{1}{f}\nu$ is $\partial^{or} \otimes \nabla$ -constant. The proposition follows now from Proposition 2.2.

3 A generalization of the Chern-Hirzebruch-Serre Lemma and applications to cohomology of Lie algebras

We generalize Lemma 3 from [2] concerning Poincaré differentiation from algebras to pairings. The assumption on finite dimensionality is superfluous.

Lemma 3.1. *Let $A_s = \bigoplus_{i=0}^n A_s^i$, $d_s : A_s \rightarrow A_s$, $s = 1, 2, 3$, be three graded differential \mathbb{R} -vector spaces such that*

- (1) $d_s[A_s^i] \subset A_s^{i+1}$,
 (2) $d_s^2 = 0$,
 (3) $d_3[A_3^{n-1}] = 0$.
 (4) $A_3^n \cong \mathbb{R}$, $A_3^i = 0$ for $i > n$.

Let

$$\cdot : A_1 \times A_2 \rightarrow A_3$$

be a pairing such that

- (5) $A_1^i \cdot A_2^j \subset A_3^{i+j}$,
 (6) $d_3(x \cdot y) = d_1x \cdot y + (-1)^{\deg x} x \cdot d_2y$,
 (7) $\cdot : A_1^r \times A_2^{n-r} \rightarrow A_3^n \cong \mathbb{R}$, $r = 0, 1, \dots, n$ are nondegenerate in the sense that the induced mappings

$$i_r : A_1^r \xrightarrow{\cong} (A_2^{n-r})^*$$

are linear isomorphisms.

Then the induced homomorphisms in cohomology

$$\cdot : H^r(A_1, d_1) \times H^{n-r}(A_2, d_2) \rightarrow H^n(A_3, d_3) \cong \mathbb{R}$$

are nondegenerate as well, i.e. the induced linear homomorphism

$$i'_r : H^r(A_1, d_1) \rightarrow (H^{n-r}(A_2, d_2))^*$$

are linear homomorphisms.

Proof. The proof is identical with the original proof by Chern-Hirzebruch-Serre for an algebra and it is sufficient to check that

$$i'_r : H^r(A_1, d_1) \xrightarrow{(i_r)_\#} H^{n-r}(A_2^*, d_2^*) \xrightarrow{\cong} H^{n-r}(A_2, d_2),$$

where (A_2^*, d_2^*) denotes the dual complex.

Now we give some applications to the cohomology of Lie algebras with coefficients. Let \mathfrak{g} be a real Lie algebra of dimension n and let

$$\nabla : \mathfrak{g} \rightarrow L_{\mathbb{R}} = \text{End } \mathbb{R} \cong \mathbb{R}$$

be an arbitrary representation in 1 dimensional vector space. We will distinguish two representations

- $\nabla_0 = 0,$
- $(\nabla_{\text{trad}})_a = \text{tr}(\text{ad}_a) \cdot \text{id}.$

We see that $\nabla_0 = \nabla_{\text{trad}}$ if and only if \mathfrak{g} is unimodular. Denote the differential with respect to ∇_{trad} by d_{trad} and the cohomology of \mathfrak{g} by $H_{\text{trad}}(\mathfrak{g})$. Straightforward computations show that $d_{\text{trad}}^{n-1} = 0$. Therefore

Proposition 3.2. $H_{\text{trad}}^n(\mathfrak{g}) = \Lambda^n \mathfrak{g}^* \cong \mathbb{R}$ for every Lie algebra.

Let us notice the following

- Remark 3.3.** (1) Each representation $\nabla : \mathfrak{g} \rightarrow L_{\mathbb{R}}$ is equal to 0 on \mathfrak{g}^2 and, conversely, each linear homomorphism $\nabla : \mathfrak{g} \rightarrow L_{\mathbb{R}}$ such that $\nabla|\mathfrak{g}^2 = 0$ is a representation.
- (2) The zero group of cohomology $H_{\nabla}^0(\mathfrak{g}) = 0$ if and only if $\nabla \neq 0$.
- (3) The multiplication of reals $\cdot : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is compatible with $(\nabla_1, \nabla_2, \nabla_3)$ if and only if $\nabla_3 = \nabla_1 + \nabla_2$.

Point (1) from the remark above implies that any linear combination of representations is a representation. Take an arbitrary representation ∇ and put

$$\nabla' = \nabla_{\text{trad}} - \nabla.$$

Then the multiplication of reals is compatible with $(\nabla', \nabla, \nabla_{\text{trad}})$ by (3) from the remark. Therefore, for differential operators $d_{\nabla'}, d_{\nabla}, d_{\text{trad}}$ the condition (6) from Lemma 3.1 holds. Since the exterior multiplication

$$\wedge : \Lambda^r \mathfrak{g}^* \times \Lambda^{n-r} \mathfrak{g}^* \rightarrow \Lambda^n \mathfrak{g}^* \cong \mathbb{R}$$

is nondegenerate then according to Lemma 3.1 the multiplication in cohomology

$$H_{\nabla'}^r(\mathfrak{g}) \times H_{\nabla'}^{n-r}(\mathfrak{g}) \rightarrow H_{\text{trad}}^n(\mathfrak{g}) \cong \mathbb{R}$$

is nondegenerate as well, i.e. in particular

$$H_{\nabla'}^0(\mathfrak{g}) \cong (H_{\nabla'}^n(\mathfrak{g}))^*.$$

Immediately from the above reasoning we obtain the following theorems.

Theorem 3.4. *The multiplication of reals is compatible with the representations $(0, \nabla_{\text{trad}}, \nabla_{\text{trad}})$ and the induced cohomology pairing*

$$H^{n-r}(\mathfrak{g}) \times H_{\text{trad}}^r(\mathfrak{g}) \rightarrow H_{\text{trad}}^n(\mathfrak{g}) \cong \mathbb{R},$$

is nondegenerate. In particular we obtain a noncanonical isomorphism

$$H_{\text{trad}}^r(\mathfrak{g}) \cong (H_{\text{trad}}^r(\mathfrak{g}))^* \cong H^{n-r}(\mathfrak{g}).$$

Theorem 3.5. ∇_{trad} *is the unique representation ∇ for which $H_{\nabla}^n(\mathfrak{g}) \neq 0$.*

Proof. For any representation ∇ take $\nabla' = \nabla_{\text{trad}} - \nabla$. By (2) from the remark above we have

$$H_{\nabla}^n(\mathfrak{g}) \cong H_{\nabla'}^0(\mathfrak{g}) \neq 0 \iff \nabla' = 0 \iff \nabla = \nabla_{\text{trad}}.$$

4 Pairings for graded filtered differential \mathbb{R} -vector spaces and spectral sequences

The aim of this chapter is to prove that for any pairing of graded regularly filtered differential \mathbb{R} -vector spaces, if the second terms of spectral sequences gives the nondegenerate pairing then the same holds for the cohomology algebras of the spaces. This holds without assumption that $\dim E_2$ is finite and generalizes the suitable theorem for graded filtered differential algebras [16].

Given three graded filtered differential \mathbb{R} -vector spaces

$$({}^r A = \bigoplus_{i \geq 0} {}^r A^i, \mathcal{d}, \mathbb{A}_j), \quad r = 1, 2, 3, \quad (8)$$

denote for shortness

$${}^r H := H({}^r A, \mathcal{d}).$$

Assume

$$\cdot : {}^1A \times {}^2A \rightarrow {}^3A$$

preserves gradations and filtrations

$${}^1A^s \cdot {}^2A^t \subset {}^3A^{s+t}, \tag{9}$$

$${}^1A_j \cdot {}^2A_k \subset {}^3A_{j+k}, \tag{10}$$

and that the differentials ${}^r d$ satisfy the compatibility condition

$${}^3d(x \cdot y) = {}^1dx \cdot y + (-1)^{\deg x} x \cdot {}^2dy. \tag{11}$$

Clearly, there exists a multiplication of cohomology classes

$$\cdot : {}^1H^j \times {}^2H^k \rightarrow {}^3H^{j+k}, \quad ([x], [y]) \mapsto [x \cdot y] ..$$

Let

$$({}^r E_s^{j,i}, {}^r d_s)$$

be spectral sequences of graded filtered differential \mathbb{R} -vector spaces (8).

Lemma 4.1. (1)

$${}^1Z_s^{j,i} \cdot {}^2Z_s^{k,l} \subset {}^3Z_s^{j+k,i+l}, \quad 0 \leq s \leq \infty,$$

(2)

$$\begin{aligned} {}^1Z_s^{j,i} \cdot {}^2D_{s-1}^{k,l} + {}^1D_{s-1}^{j,k} \cdot {}^2Z_s^{i,l} &\subset {}^3Z_{s-1}^{j+k+1,i+l-1} + {}^3D_{s-1}^{j+k,i+l}, \quad 0 \leq s < \infty, \\ {}^1Z_\infty^{j,i} \cdot {}^2D_\infty^{k,l} + {}^1D_\infty^{j,i} \cdot {}^2Z_\infty^{k,l} &\subset {}^3D_\infty^{j+k,i+l}, \quad (s = \infty). \end{aligned}$$

Proof. Straightforward calculations.

Conclusion 4.2. *There exists a multiplication of s -terms of spectral sequences*

$${}^1E_s^{j,i} \times {}^2E_s^{k,l} \rightarrow {}^3E_s^{j+k,i+l}, \quad ([x], [y]) \mapsto [x \cdot y], \quad 0 \leq s \leq \infty.$$

The differentials ${}^1d_s, {}^2d_s, {}^3d_s$ fulfils the compatibility condition with respect to the total gradation

$${}^3d_s(x \cdot y) = {}^1d_s x \cdot y + (-1)^{\text{total deg } x} x \cdot {}^2d_s y.$$

There exists a multiplication of cohomology classes of s -terms

$$H^{j,i}({}^1E_s, {}^1d_s) \times H^{k,l}({}^2E_s, {}^2d_s) \rightarrow H^{j+k,i+l}({}^3E_s, {}^3d_s), \quad ([\tilde{x}], [\tilde{y}]) \mapsto [\tilde{x} \cdot \tilde{y}].$$

The linear isomorphisms of bigraded spaces

$$\begin{aligned} {}^r\sigma_s : {}^rE_{s+1} &\rightarrow H({}^rE_s, {}^r d_s), \\ {}^r\sigma_\infty : {}^rE_\infty &\rightarrow E_0({}^rH) \end{aligned}$$

conserve the multiplications

$$\begin{aligned} {}^3\sigma_s([x] \cdot [y]) &= {}^1\sigma_s[x] \cdot {}^2\sigma_s[y], \\ {}^3\sigma_\infty(\bar{x} \cdot \bar{y}) &= {}^1\sigma_\infty(\bar{x}) \cdot {}^2\sigma_\infty(\bar{y}). \end{aligned}$$

Remark 4.3. For $s \geq i + 2$ we consider the canonical epimorphisms

$${}^r\gamma_s^{j,i} : {}^rE_s^{j,i} \cong {}^rZ_\infty^{j,i} / ({}^rZ_\infty^{j+1,i-1} + {}^rD_{s-1}^{j,i}) \twoheadrightarrow {}^rZ_\infty^{j,i} / ({}^rZ_\infty^{j+1,i-1} + {}^rD_\infty^{j,i}) \cong {}^rE_\infty^{j,i}.$$

For $s \geq i + l + 2$ the canonical epimorphisms ${}^1\gamma_s^{j,i}$, ${}^2\gamma_s^{k,l}$, ${}^3\gamma_s^{j+k,i+l}$ are compatible with multiplications

$${}^3\gamma_s^{j+k,i+l}([x] \cdot [y]) = {}^1\gamma_s^{j,i}[x] \cdot {}^2\gamma_s^{k,l}[y].$$

This implies that if spectral sequences $({}^rE_s, {}^r d_s)$ collapse at the \bar{m} th term then the canonical isomorphisms ${}^r\beta_{\bar{m}} : {}^rE_{\bar{m}} \xrightarrow{\cong} {}^rE_\infty$, see [7, Vol.III. §1.1.2], conserve bigradations and are compatible with multiplications. We recall the construction of ${}^r\beta_{\bar{m}}$. For arbitrary (j, i) we select arbitrary $s \geq \max(\bar{m}, i + 2)$ and put

$${}^r\beta_{\bar{m}}^{j,i} : {}^rE_{\bar{m}}^{j,i} \xleftarrow[\cong]{{}^r\sigma_{\bar{m}}^{j,i}} {}^rE_{\bar{m}+1}^{j,i} \xleftarrow[\cong]{} \dots \xleftarrow[\cong]{} {}^rE_s^{j,i} \xrightarrow[\cong]{{}^r\gamma_s^{j,i}} {}^rE_\infty^{j,i}.$$

The following main result of this chapter generalizes Corollary 12 from [16].

Theorem 4.4. *Given three graded filtered differential \mathbb{R} -vector spaces (8) and a pairing $\cdot : {}^1A \times {}^2A \rightarrow {}^3A$ satisfying (9), (10), (11), assume that the filtrations are regular in the sense ${}^rA_0 = {}^rA$ and that the second terms ${}^rE_2^{j,i}$ live in the rectangular $0 \leq j \leq m$, $0 \leq i \leq n$ and that ${}^3E_2^{(m+n)} = {}^3E_2^{m,n} \cong \mathbb{R}$.*

If the multiplication in the second terms

$$\langle \cdot, \cdot \rangle_2 : {}^1E_2^{(j)} \times {}^2E_2^{(m+n-j)} \rightarrow {}^3E_2^{m,n} \cong \mathbb{R}$$

is nondegenerate in the sense that

$${}^1E_2^{(j)} \xrightarrow{\cong} ({}^2E_2^{(m+n-j)})^*, \quad x \mapsto \langle x, \cdot \rangle_2,$$

is a linear isomorphism, then

- (a) ${}^3H^{m+n} \cong \mathbb{R}$,
- (b) ${}^rH^t = 0$ for $t > m + n$,
- (c) *the multiplication in cohomology classes*

$$\langle \cdot, \cdot \rangle_H : {}^1H^j \times {}^2H^{m+n-j} \rightarrow {}^3H^{m+n} \cong \mathbb{R} \tag{12}$$

is nondegenerate as well, i.e.

$${}^1H^j \xrightarrow{\cong} ({}^2H^{m+n-j})^*, \quad [x] \mapsto \langle [x], \cdot \rangle_H,$$

is a linear isomorphism.

Proof. The terms ${}^rE_3, {}^rE_4, \dots, {}^rE_\infty$ live also in the same rectangular $0 \leq j \leq m, 0 \leq i \leq n$. The bidegree argument of the second differential operator 3d_2 implies (compare with [16]) the condition ${}^3d_2[{}^3E_2^{(m+n-1)}] = 0$. By the generalized Chern-Hirzebruch-Serre Lemma 3.1 we get ${}^3E_2^{(m+n)} = {}^3E_2^{m,n} \cong \mathbb{R}$ and nondegeneracy of the multiplication for third terms. Proceeding inductively we get the same for all finite terms. The bidegree argument for the further differential operators ${}^r d_s$ implies the collapsing of spectral sequences $({}^rE_s, {}^r d_s)$, say at ${}^r m > \max(m+1, n+2)$ places. Then ${}^3E_\infty^{(m+n)} = {}^3E_\infty^{m,n} \cong \mathbb{R}$ so (a) holds because ${}^3H^{m+n} \cong {}^3E_\infty^{(m+n)} \cong \mathbb{R}$ and next, for $\bar{m} \geq \max({}^1m, {}^2m, {}^3m)$ the canonical isomorphisms ${}^r\beta_{\bar{m}}$ (see Remark 4.3) are compatible with multiplications. In consequence, the multiplication in the infinite terms

$$\cdot : {}^1E_\infty^{(j)} \times {}^2E_\infty^{(m+n-j)} \rightarrow {}^3E_\infty^{m,n} \cong \mathbb{R} \tag{13}$$

is nondegenerate as well.

It remains to prove the nondegeneracy of the multiplication of cohomology classes (12). The spaces rH possess a natural graded filtration ${}^rH^{j,i}$, and thanks to the regularity of filtrations we have

$${}^rH^t = {}^rH^{0,t} \supset {}^rH^{1,t-1} \supset \dots \supset {}^rH^{t,0} \supset 0 \tag{14}$$

and a noncanonical isomorphism

$${}^rH^t \cong ({}^rH^{0,t}/{}^rH^{1,t-1}) \oplus ({}^rH^{1,t-1}/{}^rH^{2,t-2}) \oplus \dots \oplus {}^rH^{t,0} = \bigoplus_{j+i=t} E_0^{j,i}({}^rH). \tag{15}$$

Analogously to the proof of Theorem 11 from [16] we assert that

$$E_0^{(m+n)}({}^rH) = E_0^{m,n}({}^rH) = {}^rH^{m,n}, \tag{16}$$

and

$${}^rH^{j,i} = {}^rH^{j+1,i-1} \text{ for } j > m \text{ or } i > n.$$

Therefore by (15) ${}^rH^t = 0$ for $t > m+n$ which proves (b). As in [16] we check the rule:

- if ${}^1\sigma_\infty^{j,i}(\bar{x}) = [x]$ for $\bar{x} \in {}^1E_\infty^{j,i}$, $x \in {}^1H^{j,i}$, and if ${}^2\sigma_\infty^{m-j,n-i}(\bar{y}) = [y]$ for $\bar{y} \in {}^2E_\infty^{m-j,n-i}$, $y \in {}^2H^{m-j,n-i}$, then

$${}^3\sigma_\infty^{m,n}(\bar{x} \cdot \bar{y}) = [x] \cdot [y] = x \cdot y. \tag{17}$$

We fix generators $\xi_\infty \in {}^3E_\infty^{m,n}$ and $\xi_H \in {}^3H^{m,n}$ in such a way that ${}^3\sigma_\infty^{m,n}(\xi_\infty) = \xi_H$. Consider the pairings, see (13),

$$\begin{aligned} \langle \cdot, \cdot \rangle_\infty : {}^1E_\infty^{(j)} \times {}^2E_\infty^{(m+n-j)} &\rightarrow \mathbb{R}, \quad \langle \bar{x}, \bar{y} \rangle_\infty \cdot \xi_\infty = \bar{x} \cdot \bar{y}, \\ \langle \cdot, \cdot \rangle_H : {}^1H^j \times {}^2H^{m+n-j} &\rightarrow \mathbb{R}, \quad \langle x, y \rangle_H \cdot \xi_H = x \cdot y. \end{aligned}$$

By (17) we have

$$\langle \bar{x}, \bar{y} \rangle_\infty = \langle x, y \rangle_H \tag{18}$$

where ${}^1\sigma_\infty^{j,i}(\bar{x}) = [x]$ and ${}^2\sigma_\infty^{m-j,n-i}(\bar{y}) = [y]$. According to (13) the pairing $\langle \cdot, \cdot \rangle_\infty$ is nondegenerate, that is ${}^1E_\infty^{(j)} \cong ({}^2E_\infty^{(m+n-j)})^*$. Consider the induced linear mapping

$$\kappa : {}^1H^j \rightarrow ({}^2H^{m+n-j})^*, \quad x \mapsto \langle x, \cdot \rangle_H.$$

Similarly to [16] we easily check the monomorphy of κ . It remains to check that κ is an epimorphism.

Take a linear function $0 \neq l : {}^2H^{m+n-j} \rightarrow \mathbb{R}$ and consider the filtration (14) for $r = 2$ and $t = m + n - j$. Let $V^p \subset {}^2H^{p,m+n-j-p}$ be a subspace complementary to ${}^2H^{p+1,m+n-j-p-1}$, $p = 0, 1, \dots, m + n - j$ and

$$\psi^p : V^p \rightarrow E_0^{p,m+n-j-p}({}^2H), \quad x \mapsto [x],$$

the induced isomorphism. Put

$$\psi = \sum_p \psi^p : {}^2H^{m+n-j} = \bigoplus_p V^p \cong \bigoplus_p E_0^{p,m+n-j-p}({}^2H).$$

The composition $l \circ \psi^{-1} \in (\bigoplus_p E_0^{p,m+n-j-p}({}^2H))^*$ determines a family of linear functions $l_0^p \in (E_0^{p,m+n-j-p}({}^2H))^*$. Define

$$I_l = \{p; l_0^p \neq 0\}.$$

For each $p \in I_l$ we define - through isomorphisms

$${}^2\sigma_\infty^{p,m+n-j-p} : {}^2E_\infty^{p,m+n-j-p} \xrightarrow{\cong} E_0^{p,m+n-j-p}({}^2H)$$

- a linear nonzero functions

$$l_\infty^p \in ({}^2E_\infty^{p,m+n-j-p})^*, \quad l_\infty^p = l_0^p \circ {}^2\sigma_\infty^{p,m+n-j-p}.$$

The nondegenerate pairing $\langle \cdot, \cdot \rangle_\infty : {}^1E_\infty^{m-p,p+j-m} \times {}^2E_\infty^{p,m+n-j-p} \rightarrow \mathbb{R}$ determines an element $0 \neq \bar{x}^{m-p} \in {}^1E_\infty^{m-p,p+j-m}$ such that $\langle \bar{x}^{m-p}, \cdot \rangle_\infty = l_\infty^p \in ({}^2E_\infty^{p,m+n-j-p})^*$. Let ${}^1\sigma_\infty^{m-p,p+j-m}(\bar{x}^{m-p}) = [x^{m-p}] \in E_0^{m-p,p+j-m}({}^1H)$, where

$$x^{m-p} \in {}^1H^{m-p,p+j-m} \quad \text{and} \quad x^{m-p} \notin {}^1H^{m-p+1,p+j-m-1}.$$

Put

$$x := \sum_{p \in I_l} x^{m-p}.$$

We prove the equality

$$\kappa(x) = \langle x, \cdot \rangle_H = l \in ({}^2H^{m+n-j})^*.$$

Since ${}^2H^{m+n-j} = \bigoplus_p V^p$, we need only to prove $\kappa(x)(y^p) = \langle x, \cdot \rangle_H(y^p) = l(y^p)$ for $y^p \in V^p \subset {}^2H^{p,m+n-j-p}$. If $p \notin I_l$, i.e. $l_0^p = 0$, then $l(y^p) = 0$ and for all $p' \in I_l$ by (16) and (17)

$$\langle x^{m-p'}, y^p \rangle_H \cdot \xi_H = x^{m-p'} \cdot y^p = [x^{m-p'}] \cdot [y^p] \in E_0^{m-p+p',n+p-p'}({}^3H) = 0.$$

If $p \in I_l$ and $0 \neq y^p \in V^p$ then

$$\begin{aligned} l(y^p) &= (l|V^p) \circ (\psi^p)^{-1} [y] = l_0^p [y^p] = l_\infty^p \circ (\sigma_\infty^{p,m+n-j-p})^{-1} ([y^p]) \\ &= \langle \bar{x}^{m-p}, (\sigma_\infty^{p,m+n-j-p})^{-1} ([y^p]) \rangle_\infty \\ &\stackrel{(18)}{=} \langle x^{m-p}, y^p \rangle_H \\ &= \langle x, y^p \rangle_H. \end{aligned}$$

The last equation holds because for $p' \neq p, p' \in I_l$, we have

$$0 \neq [x^{m-p'}] \in E_0^{m-p',p'+j-m} ({}^1H)$$

and by (16) and (17)

$$x^{m-p'} \cdot y^p = [x^{m-p'}] \cdot [y^p] \in E_0^{m-p',p'+j-m} ({}^3H) = 0.$$

5 Hochschild-Serre filtration and the spectral sequence for transitive Lie algebroids

We fix a transitive Lie algebroid $A = (A, [\cdot, \cdot], \#_A)$ with the Atiyah sequence $0 \rightarrow \mathfrak{g} \hookrightarrow A \xrightarrow{\#_A} TM \rightarrow 0$ and a representation $\nabla : A \rightarrow A(\xi)$ of a Lie algebroid A on a vector bundle ξ . ∇ is a homomorphism of Lie algebroids, then ∇ induces a homomorphism of vector bundles $\nabla^+ : \mathfrak{g} \rightarrow \text{End}(\xi)$

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{\nabla^+} & \text{End}(\xi) \\ \downarrow & & \downarrow \\ A & \xrightarrow{\nabla} & A(\xi) \end{array}$$

and $\nabla_x^+ : \mathfrak{g}_x \rightarrow \text{End}(\xi_x)$ is a representation of the isotropy Lie algebra \mathfrak{g}_x in the vector space ξ_x . We will consider the pair of \mathbb{R} -Lie algebras $(\mathfrak{g}, \mathfrak{k})$ where

$$\mathfrak{g} = \Gamma(A), \quad \mathfrak{k} = \Gamma(\mathfrak{g}).$$

Below, the elements of \mathfrak{g} will be denoted by $\gamma, \gamma_1, \gamma_2, \dots$ while elements of \mathfrak{k} by $\sigma, \sigma_1, \sigma_2, \dots$. Of course, \mathfrak{k} is an ideal of \mathfrak{g} (actually, \mathfrak{k} is $C^\infty(M)$ -Lie algebra but it is not interesting here). The space $\Gamma(\xi)$ is a \mathfrak{g} -modul with respect to the induced representation denoted by the same letter $\nabla : \Gamma(A) \rightarrow \Gamma(A(\xi)) \subset L_{\Gamma(\xi)}$. Following Hochschild-Serre [9] we can consider a graded cochain group of \mathbb{R} -linear alternating functions

$$A_{\mathbb{R}} = \bigoplus_{i \geq 0} A^i, \quad A^i = C^i(\mathfrak{g}, \Gamma(\xi)),$$

with the \mathbb{R} -differential operator of degree 1

$$d_{\nabla} : C^i(\mathfrak{g}, \Gamma(\xi)) \rightarrow C^{i+1}(\mathfrak{g}, \Gamma(\xi))$$

defined by the standard formula

$$(d_{\nabla} f)(\gamma_0, \dots, \gamma_t) = \sum_i (-1)^i \nabla_{\gamma_i} (f(\gamma_0, \dots, \hat{\gamma}_i, \dots, \gamma_t)) + \sum_{i < j} (-1)^{i+j} f([\gamma_i, \gamma_j], \dots, \hat{\gamma}_i, \dots, \hat{\gamma}_j, \dots).$$

For the trivial representation $\partial_A : A \rightarrow A(M \times \mathbb{R})$, $(\partial_A)_\gamma (f) = \partial_{\#(\gamma)} (f)$, this operator is denoted by d_A . Clearly, for a real alternating t -cocycle φ and $\sigma \in \Gamma(\xi)$ we get

$$d_{\nabla}(\varphi \otimes \sigma) = d_A \varphi \otimes \sigma + (-1)^t \varphi \wedge d_{\nabla} \sigma.$$

In the space $\bigoplus_{t \geq 0} A^t$ we have the Hochschild-Serre filtration $A_j \subset A_{\mathbb{R}}$ as follows: $A_j = A_{\mathbb{R}}$ for $j \leq 0$. If $j > 0$, $A_j = \bigoplus_{t \geq j} A_j^t$, $A_j^t = A_j \cap A^t$, where A_j^t consists of all those t -cochains f for which $f(\gamma_1, \dots, \gamma_t) = 0$ whenever $t - j + 1$ of the arguments γ_i belongs to \mathfrak{k} . In this way we have obtained a graded filtered differential \mathbb{R} -vector space

$$(A_{\mathbb{R}} = \bigoplus_{t \geq 0} A^t, d_{\nabla}, A_j) \quad (19)$$

and we can use its spectral sequence

$$(E_s^{j,i}, d_s). \quad (20)$$

Following K. Mackenzie [17] (see also V.Itskov, M.Karashev, and Y.Vorobjev [11]) we will consider the $C^\infty(M)$ -submodule of $C^\infty(M)$ -linear alternating cochains with values in the vector bundle ξ (i.e. A -differential ξ -valued forms)

$$\Omega^t(A, \xi) \subset C^t(\mathfrak{g}, \Gamma(\xi))$$

and the induced filtration

$$\Omega_j = \Omega_j(A, \xi) = A_j \cap \Omega(A, \xi)$$

of $C^\infty(M)$ -modules. The differential d_{∇} of a $C^\infty(M)$ -cochain is a $C^\infty(M)$ -cochain, so we get $d_{A, \nabla} : \Omega(A, \xi) \rightarrow \Omega(A, \xi)$. We obtain in this way a graded filtered differential space

$$(\Omega(A, \xi) = \bigoplus_t \Omega^t(A, \xi), d_{A, \nabla}, \Omega_j) \quad (21)$$

and its spectral sequence

$$(E_{A,s}^{j,i}, d_{A, \nabla, s}). \quad (22)$$

Now we consider as well a submodule of $C^\infty(M)$ -linear alternating cochains with compact support $\Omega_c^t(A, \xi) \subset \Omega^t(A, \xi)$ and the corresponding filtration

$$\Omega_{c,j} = \Omega_j \cap \Omega_c(A, \xi)$$

of $C^\infty(M)$ -modules. Since $\text{supp } d_{\nabla} f \subset \text{supp } f$ then we obtain

$$d_{A_c, \nabla} : \Omega_c(A, \xi) \rightarrow \Omega_c(A, \xi)$$

and we get a graded filtered differential space with compact support

$$(\Omega_c(A, \xi) = \bigoplus_t \Omega_c^t(A, \xi), d_{A_c, \nabla}, \Omega_{c,j}) \tag{23}$$

and its spectral sequence

$$(E_{A_c, s}^{j,i}, d_{A_c, \nabla, s}). \tag{24}$$

Sometimes we can deduce directly properties of the last two spectral sequences (22), (24) from the suitable properties of (20), see [9], denoted further by $\boxed{A_{\mathbb{R}}}$, sometimes we must use some additional observations.

Lemma 5.1. *The homomorphisms ρ_0 and $\rho_{c,0}$ in the sequence*

$$E_{A_c, 0}^{j,i} = \Omega_{c,j}^{j+i} / \Omega_{c,j+1}^{j+i} \xrightarrow{\rho_{c,0}} \Omega_j^{j+i} / \Omega_{j+1}^{j+i} = E_{A,0}^{j,i} \xrightarrow{\rho_0} A_j^{j+i} / A_{j+1}^{j+i} = E_0^{j,i}$$

are monomorphisms. For differentials $d_{A_c, \nabla}, d_{A, \nabla}, d_0$ the following diagram is commutative

$$\begin{array}{ccccc} E_{A_c, 0}^{j,i} & \xrightarrow{\rho_{c,0}} & E_{A, 0}^{j,i} & \xrightarrow{\rho_0} & E_0^{j,i} \\ \downarrow d_{A_c, \nabla, 0}^{j,i} & & \downarrow d_{A, \nabla, 0}^{j,i} & & \downarrow d_0^{j,i} \\ E_{A_c, 0}^{j,i+1} & \xrightarrow{\rho_{c,0}} & E_{A, 0}^{j,i+1} & \xrightarrow{\rho_0} & E_0^{j,i+1} \end{array}$$

From

- $\boxed{A_{\mathbb{R}}}$ For R -cochains there exists an isomorphism

$$a^{j,i} : E_0^{j,i} \rightarrow C^j(\mathfrak{g}/\mathfrak{k}, C^i(\mathfrak{k}, \Gamma(\xi)))$$

such that

$$a^{j,i}[f]([\gamma_1], \dots, [\gamma_j])(\sigma_1, \dots, \sigma_i) = f(\sigma_1, \dots, \sigma_i, \gamma_1, \dots, \gamma_j), \tag{25}$$

we can easily obtain the following

Conclusion 5.2. *The homomorphisms*

$$\begin{aligned} \alpha_A^{j,i} &: E_{A,0}^{j,i} \rightarrow \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi) \\ \alpha_{A_c}^{j,i} &: E_{A_c,0}^{j,i} \rightarrow \Omega_c^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi) \end{aligned}$$

defined by the formula

$$\alpha_A^{j,i}[f](X_1, \dots, X_j)(\sigma_1, \dots, \sigma_i) = f(\sigma_1, \dots, \sigma_i, \lambda X_1, \dots, \lambda X_j),$$

$X_{j'} \in \mathfrak{X}(M), \sigma_{i'} \in \mathfrak{k}$, ($\alpha_{A_c}^{j,i}$ defined by the identical formula) where $\lambda : TM \rightarrow A$ is an arbitrary connection, are correctly defined linear isomorphisms of $C^\infty(M)$ -modules.

Proof. Monomorphy of $\alpha_A^{j,i}$ and $\alpha_{A_c}^{j,i}$ follows from the commutativity of the diagram

$$\begin{array}{ccccc}
 E_{A_c,0}^{j,i} & \xrightarrow{\rho_{c,0}} & E_{A,0}^{j,i} & \xrightarrow{\rho_0} & E_0^{j,i} \\
 \downarrow a_{A_c}^{j,i} & & \downarrow a_A^{j,i} & & \downarrow a^{j,i} \\
 \Omega_c^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi) & \xrightarrow{\bar{\rho}_{c,0}} & \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi) & \xrightarrow{\bar{\rho}_0} & C^j(\mathfrak{g}/\mathfrak{k}, C^i(\mathfrak{k}, \Gamma(\xi))).
 \end{array}$$

To prove that $a_A^{j,i}$ is an epimorphism it is sufficient to check that if $a^{j,i}[f]$ is a $C^\infty(M)$ -linear cochain, i.e. $a^{j,i}[f] = \bar{\rho}_0(\bar{f})$ for some $\bar{f} \in \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi)$, i.e.

$$\bar{f}(\#_A(\gamma_1), \dots, \#_A(\gamma_j))(\sigma_1, \dots, \sigma_i) = f(\sigma_1, \dots, \sigma_i, \gamma_1, \dots, \gamma_j),$$

then there exists a representative $f' \in [f] \in E_{A,0}^{j,i}$ which is $C^\infty(M)$ -linear cochain such that $a_A^{j,i}[f'] = \bar{f}$. To this end take a connection form $\omega_0 : A \rightarrow \mathfrak{g}$ corresponding to λ and put $f'(\gamma'_1, \dots, \gamma'_j, \gamma_1, \dots, \gamma_i) = f(\omega_0(\gamma'_1), \dots, \omega_0(\gamma'_j), \gamma_1, \dots, \gamma_i)$. Then f' fulfils the desired conditions.

- $\square_{\mathbb{R}}$ Through isomorphism $a^{j,i}$ the differential $d_0^{j,i}$ becomes a differentiation of values with respect to the differential

$$d_{\nabla_{ol}} : C^i(\mathfrak{k}, \Gamma(\xi)) \rightarrow C^{i+1}(\mathfrak{k}, \Gamma(\xi)) \tag{26}$$

($\iota : \mathfrak{k} \hookrightarrow \mathfrak{g}$, is the inclusion),

$$\begin{aligned}
 \tilde{d}_{\nabla_{ol}} : C^j(\mathfrak{g}/\mathfrak{k}, C^i(\mathfrak{k}, \Gamma(\xi))) &\rightarrow C^j(\mathfrak{g}/\mathfrak{k}, C^{i+1}(\mathfrak{k}, \Gamma(\xi))), \\
 \tilde{d}_{\nabla_{ol}}(f)([\gamma_1], \dots, [\gamma_j]) &= d_{\nabla_{ol}}(f([\gamma_1], \dots, [\gamma_j])).
 \end{aligned}$$

In conclusion, the differentials $d_{A,\nabla,0}^{j,i}$ and $d_{A_c,\nabla,0}^{j,i}$ becomes (through the isomorphisms $a_A^{j,i}$ and $a_{A_c}^{j,i}$) differentials of values with respect to

$$d_{\nabla+} : \Lambda^i \mathfrak{g}^* \otimes \xi \rightarrow \Lambda^{i+1} \mathfrak{g}^* \otimes \xi,$$

namely

$$\begin{aligned}
 \tilde{d}_{\nabla+} : \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi) &\rightarrow \Omega^j(M, \Lambda^{i+1} \mathfrak{g}^* \otimes \xi), \\
 \tilde{d}_{\nabla+}(f)(X_1, \dots, X_j) &= d_{\nabla+}(f(X_1, \dots, X_j)).
 \end{aligned}$$

Analogously we obtain a differential $\tilde{d}_{c,\nabla+}$ for compact supports.

Remark 5.3. According to K.Mackenzie [17, Th.2.5, p.201] the homomorphisms $d_{\nabla+}^i : \Lambda^i \mathfrak{g}^* \otimes \xi \rightarrow \Lambda^{i+1} \mathfrak{g}^* \otimes \xi$ are locally of constant rank, and consequently, there are well-defined vector bundles $Z^i = \ker d_{\nabla+}^i$, $B^i = \text{Im } d_{\nabla+}^{i-1}$ and $H^i(\mathfrak{g}, \xi) = Z^i/B^i$ such that $\Gamma(H^i(\mathfrak{g}, \xi)) = H^i(\Gamma(\Lambda \mathfrak{g}^* \otimes \xi), d_{\nabla+})$. Clearly,

$$H^i(\mathfrak{g}, \xi)_x = H^i(\Lambda \mathfrak{g}_x^* \otimes \xi_x, d_{\nabla+^x}).$$

Therefore

$$\begin{aligned}
 H(\Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi), \tilde{d}_{\nabla+}) &\cong \Omega^j(M, H_{\nabla+}^i(\mathfrak{g}, \xi)), \\
 H(\Omega_c^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi), \tilde{d}_{c,\nabla+}) &\cong \Omega_c^j(M, H_{\nabla+}^i(\mathfrak{g}, \xi)).
 \end{aligned}$$

From the above we obtain isomorphisms of $C^\infty(M)$ -modules

$$\begin{aligned} (a_A^{j,i})_\# &: H^i(E_{A,0}^{j,*}, d_{A,\nabla,0}^{j,*}) \xrightarrow{\cong} \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi), \\ (a_{A_c}^{j,i})_\# &: H^i(E_{A_c,0}^{j,*}, d_{A_c,\nabla,0}^{j,*}) \xrightarrow{\cong} \Omega_c^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi). \end{aligned}$$

Now we pass to consideration of the modules Z_s, D_s, E_s and $Z_{A,s}, D_{A,s}, E_{A,s}$ and $Z_{A_c,s}, D_{A_c,s}, E_{A_c,s}$ for three spectral sequences for graded, filtered, differential spaces (19), (21), (23), respectively. Immediately by definitions we get

Lemma 5.4. (1) $Z_{A,s}^j = Z_s^j \cap \Omega(A, \xi)$, (2) $D_{A,s}^j = D_s^j \cap \Omega(A, \xi)$, (3) $Z_{A_c,s}^j = Z_{A,s}^j \cap \Omega_c(A, \xi)$.

Fix an auxiliary connection $\lambda : TM \rightarrow A$ and for $f \in Z_{A,1}^{j,i} \subset \Omega^{j+i}(A, \xi)$ we define

$$\bar{f}_j \in \Omega^j(M, \Lambda^i \mathfrak{g}^* \otimes \xi)$$

by the formula

$$\begin{aligned} \bar{f}_j(X_1, \dots, X_j)(\sigma_1, \dots, \sigma_i) &= f(\lambda X_1, \dots, \lambda X_j, \sigma_1, \dots, \sigma_i) \\ &= (-1)^{ji} f(\sigma_1, \dots, \sigma_i, \lambda X_1, \dots, \lambda X_j). \end{aligned}$$

Lemma 5.5. If $f \in Z_{A,1}^{j,i}$ then $\bar{f}_j(X_1, \dots, X_j) \in \Gamma(\Lambda^i \mathfrak{g}^* \otimes \xi)$ is a $d_{\nabla+}$ -cocycle independent of the choice of λ .

Proof. For $f \in Z_{A,1}^{j,i} \subset \Omega_j^{j+i}(A, \xi) \subset A_s^{j+i}$ we take a cochain $f_j \in C^j(\mathfrak{g}, C^i(\mathfrak{g}, \Gamma(\xi)))$ defined by $f_j(\gamma_1, \dots, \gamma_j)(\gamma'_1, \dots, \gamma'_i) = f(\gamma_1, \dots, \gamma_j, \gamma'_1, \dots, \gamma'_i)$, see [9]. From the equalities

$$Z_{A,1}^{j,i} = \{f \in \Omega_j^{j+i}; d_{\nabla} f \in \Omega_{j+1}^{j+i+1}\} = Z_1^{j,i} \cap \Omega^{j+i}(A, \xi)$$

we get (see [H-S]) that $\iota_j^*(f_j(\gamma_1, \dots, \gamma_j)) \in C^j(\mathfrak{k}, \Gamma(\xi))$, where

$$\iota_j^* : C^j(\mathfrak{g}, \Gamma(\xi)) \rightarrow C^j(\mathfrak{k}, \Gamma(\xi)), \quad \iota_j^*(g) = g|_{\mathfrak{k} \times \dots \times \mathfrak{k}}$$

is a $(\nabla \circ \iota : \mathfrak{k} \rightarrow L_{\Gamma(\xi)})$ -cocycle and that this cocycle depends only on the equivalence class $[\gamma_{j'}] \in \mathfrak{g}/\mathfrak{k} \cong \mathfrak{X}(M)$, i.e. on the anchors of the elements $\gamma_{j'}$, i.e. on $\#_A(\gamma_{j'})$. But $\iota_j^*(f_j(\gamma_1, \dots, \gamma_j))$ is $C^\infty(M)$ -linear $\iota_j^*(f_j(\gamma_1, \dots, \gamma_j)) \in \Gamma(\Lambda^i \mathfrak{g}^* \otimes \xi)$ therefore the condition $d_{\nabla \circ \iota}(\iota_j^*(f_j(\gamma_1, \dots, \gamma_j))) = 0$ is equivalent to

$$d_{\nabla+}(\iota_j^*(f_j(\gamma_1, \dots, \gamma_j))) = 0.$$

The equality $\iota_j^*(f_j(\gamma_1, \dots, \gamma_j)) = \bar{f}_j(\#_A(\gamma_1), \dots, \#_A(\gamma_j))$ proves the lemma.

We recall that

$$\begin{aligned} E_{A,1}^{j,i} &= Z_{A,1}^{j,i} / (Z_{A,0}^{j+1,i-1} + D_{A,0}^{j,i}) \\ &= \{f \in \Omega_j^{j+i}(A, \xi); d_{\nabla} f \in \Omega_{j+1}^{j+i+1}\} / (\Omega_{j+1}^{j+i} + d_{\nabla}[\Omega_j^{j+i-1}]) \end{aligned}$$

and analogously for $E_{A_c,1}^{j,i}$.

Lemma 5.6. *The homomorphisms*

$$\begin{aligned} \Psi_{A,1} &: E_{A,1}^{j,i} \rightarrow \Omega^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi)), \quad [f] \mapsto (-1)^{ji} [\bar{f}_j], \\ \Psi_{A_c,1} &: E_{A_c,1}^{j,i} \rightarrow \Omega_c^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi)), \quad [f] \mapsto (-1)^{ji} [\bar{f}_j], \end{aligned}$$

are isomorphisms of $C^\infty(M)$ -modules.

Proof. Clearly, we need to notice only that $\Psi_{A,1}$ is a composition of isomorphisms

$$\Psi_{A,1} : E_{A,1}^{j,i} \xrightarrow[\cong]{\sigma_{A,0}^{j,i}} H^i(E_{A,0}^{j,*}, d_{A,0}^{j,*}) \xrightarrow[\cong]{(a_A^{j,i})^\#} \Omega^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi))$$

and analogously for compact supports.

From the above lemmas we see that the canonical homomorphism $E_{A_c,1}^{j,i} \rightarrow E_{A,1}^{j,i}$ is a monomorphism.

- $\boxed{A_{\mathbb{R}}}$ *There exists a representation [precisely, a Lie derivation]*

$$\mathcal{L}^i : \mathfrak{g} \rightarrow L_{C^i(\mathfrak{k}, \Gamma(\xi))}$$

defined by the formula

$$(\mathcal{L}_\gamma^i f)(\sigma_1, \dots, \sigma_i) = \nabla_\gamma(f(\sigma_1, \dots, \sigma_i)) - \sum_t f(\sigma_1, \dots, [\gamma, \sigma_t], \dots, \sigma_i).$$

\mathcal{L}_γ^i commutes with \mathbb{R} -differential operator $d_{\nabla_{ol}}$, see (26), induces a representation in cohomology

$$\mathcal{L}^{\#,i} : \mathfrak{g} \rightarrow L_{H_{\nabla_{ol}}^i(\mathfrak{k}, \Gamma(\xi))}$$

and $\mathfrak{k} \subset \ker \mathcal{L}^{\#,i}$ (because $\mathcal{L}_\sigma^i f = d_{\nabla_{ol}}(\iota_\sigma f)$ if f is a $d_{\nabla_{ol}}$ -cocycle.). It produces a representation

$$[\mathcal{L}^{\#,i}] : \mathfrak{g}/\mathfrak{k} \rightarrow L_{H_{\nabla_{ol}}^i(\mathfrak{k}, \Gamma(\xi))}.$$

Noticing that a Lie derivation of a $C^\infty(M)$ -linear cochain is $C^\infty(M)$ -linear too, we can pass to $\Gamma(\Lambda^i \mathfrak{g}^* \otimes \xi)$. Additionally we observe that $\mathcal{L}_\gamma^i : \Gamma(\Lambda^i \mathfrak{g}^* \otimes \xi) \rightarrow \Gamma(\Lambda^i \mathfrak{g}^* \otimes \xi)$ is a covariant derivative operator with the anchor $\#_A(\gamma)$ and \mathcal{L}_γ^i is $C^\infty(M)$ -linear with respect to γ . In conclusion we obtain a representation of the Lie algebroid A in the vector bundle $\Lambda^i \mathfrak{g}^* \otimes \xi$

$$\mathcal{L}_A^i : A \rightarrow A(\Lambda^i \mathfrak{g}^* \otimes \xi).$$

Lemma 5.7. *The representation \mathcal{L}_A^i coincides with the adjoint representation of A in $\Lambda^i \mathfrak{g}^*$ cross ∇ , $\mathcal{L}_A^i = ad_A \otimes \nabla$.*

Proof. The adjoint representation $ad_A : A \rightarrow A(\mathfrak{g})$, $ad_A(\gamma)(\sigma) = [\gamma, \sigma]$, induces the one in the associated bundle $\Lambda^i \mathfrak{g}^*$ (denoted also ad_A) and its tensor product with ∇ is just equal to \mathcal{L}_A^i .

The representation \mathcal{L}_A^i induces the one in cohomology

$$\mathcal{L}_A^{\#,i} : A \rightarrow A (H_{\nabla^+}^i(\mathfrak{g}, \xi))$$

such that $\mathfrak{g} \subset \ker \mathcal{L}_A^{\#,i}$ (indeed, $(\mathcal{L}_A^i)_\sigma(f) = d_{\nabla^+}(\iota_\sigma f)$ for a d_{∇^+} -cocycle f). Therefore, we obtain a flat covariant derivative

$$\nabla^i : TM \rightarrow A (H_{\nabla^+}^i(\mathfrak{g}, \xi)) \tag{27}$$

by the formula

$$\nabla_X^i([f]) = (\mathcal{L}_A^{\#,i})_{\lambda X}([f]) = [(\mathcal{L}_A^i)_{\lambda X}(f)]$$

where for a d_{∇^+} -cocycle $f \in \Lambda^i \mathfrak{g}^* \otimes \xi$

$$(\mathcal{L}_A^i)_{\lambda X}(f)(\sigma_1, \dots, \sigma_i) = \nabla_{\lambda X}(f(\sigma_1, \dots, \sigma_i)) - \sum_t f(\sigma_1, \dots, [\lambda X, \sigma_t], \dots, \sigma_i).$$

Remark 5.8. For the trivial representation $\nabla = \partial_A$ we get a flat structure in the cohomology bundle $H^i(\mathfrak{g})$. If the structure Lie algebras \mathfrak{g}_x are unimodular then $H^n(\mathfrak{g}) = \Lambda^n \mathfrak{g}^*$ and the induced flat covariant derivative $\partial_A^n : TM \rightarrow A(\Lambda^n \mathfrak{g}^*)$ is defined by

$$((\partial_A^n)_X f)(\sigma_1, \dots, \sigma_n) = X(f(\sigma_1, \dots, \sigma_n)) - \sum_i f(\sigma_1, \dots, [\lambda X, \sigma_i], \dots, \sigma_n).$$

This flat structure coincides with the flat structure in $\Lambda^n \mathfrak{g}^*$ defined in the paper [16] via some system $\mathcal{A} = \{\tilde{\varphi}_U^n\}$ of local trivializations with locally-constant transitive functions. We recall that $\tilde{\varphi}_U^n : U \times \Lambda^n \mathfrak{g}^* \rightarrow \Lambda^n \mathfrak{g}^*$ (\mathfrak{g} is the typical fiber of \mathfrak{g}) is determined by a local trivialization $\varphi_U : A_U \rightarrow TU \times \mathfrak{g}$ of the Lie algebroid A in the following way: φ_U induces a local trivialization $\varphi_U^+ : \mathfrak{g}_U \rightarrow U \times \mathfrak{g}$ of the adjoint Lie Algebra Bundle \mathfrak{g} and we put $(\tilde{\varphi}_U^n)_x = \Lambda^n (\varphi_U^+)_x^*$.

Now, we carry over the differentials $d_{A,1}^{j,i} : E_{A,1}^{j,i} \rightarrow E_{A,1}^{j+1,i}$, $d_{A_c,1}^{j,i} : E_{A_c,1}^{j,i} \rightarrow E_{A_c,1}^{j+1,i}$, to the spaces $\Omega^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi))$ and $\Omega_c^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi))$, respectively, via the isomorphisms $\Psi_{A,1}$ and $\Psi_{A_c,1}$. Since the canonical homomorphism

$$\Gamma(H_{\nabla^+}^i(\mathfrak{g}, \xi)) \rightarrow H_{\nabla^{ol}}^i(\mathfrak{k}, \Gamma(\xi))$$

is not a monomorphism unless the Lie algebra bundle \mathfrak{g} is trivial, we can not infer the form of this differentials immediately from the level of \mathbb{R} -cochains and its spectral sequence (20). In comparing of the cohomology classes from $H_{\nabla^{ol}}^i(\mathfrak{k}, \Gamma(\xi))$ having representative of $C^\infty(M)$ -linear cochains we must see whether these representatives differ by a $C^\infty(M)$ -linear cochain.

Proposition 5.9. *The following diagrams are commutative*

$$\begin{array}{ccc} E_{A,1}^{j,i} & \xrightarrow{d_{A,1}^{j,i}} & E_{A,1}^{j+1,i} \\ \downarrow \Psi_{A,1}^{j,i} & & \downarrow \Psi_{A,1}^{j+1,i} \\ \Omega^j(M, H_{\nabla^+}^i(\mathfrak{g}, \xi)) & \xrightarrow{(-1)^i d_{\nabla^+}^i} & \Omega^{j+1}(M, H_{\nabla^+}^i(\mathfrak{g}, \xi)) \end{array}$$

$$\begin{array}{ccc}
E_{A_c,1}^{j,i} & \xrightarrow{d_{A_c,1}^{j,i}} & E_{A_c,1}^{j+1,i} \\
\downarrow \Psi_{A_c,1}^{j,i} & & \downarrow \Psi_{A_c,1}^{j+1,i} \\
\Omega_c^j(M, H_{\nabla^+}^i(\mathbf{g}, \xi)) & \xrightarrow{(-1)^i d_{\nabla^+}^i} & \Omega_c^{j+1}(M, H_{\nabla^+}^i(\mathbf{g}, \xi))
\end{array}$$

Proof. The calculations identical as in the \mathbb{R} -linear cochains [9] yield for $f \in Z_{A,1}^{j,i} \subset Z_1^{j,i}$ the following formulae

$$(-1)^i (d_{\nabla^+}^i \circ \Psi_{A,1}^{j,i} [f]) (X_1, \dots, X_{j+1}) = (-1)^{(j+1)i} [\rho_f (X_1, \dots, X_{j+1})],$$

$$\begin{aligned}
& (\Psi_{A,1}^{j+1,i} \circ d_{A,1}^{j,i} [f]) (X_1, \dots, X_{j+1}) \\
&= (-1)^{(j+1)i} [\rho_f (X_1, \dots, X_{j+1}) - d_{\nabla^+} ((-1)^j \bar{f}_{j+1} (X_1, \dots, X_{j+1}))]
\end{aligned}$$

where $\rho_f \in \Omega^{j+1}(M, \Lambda^i \mathbf{g}^* \otimes \xi)$ and $\rho_f (X_1, \dots, X_{j+1})$ is a d_{∇^+} -cocycle defined by

$$\begin{aligned}
& \rho_f (X_1, \dots, X_{j+1}) (\sigma_1, \dots, \sigma_i) \\
&= \sum_{t=1}^{j+1} (-1)^{t+1} \nabla_{\lambda X_t} (\bar{f}_j (X_1, \dots, \hat{t} \dots, X_{j+1}) (\sigma_1, \dots, \sigma_i)) + \\
&+ \sum_{t=1}^{j+1} (-1)^t \sum_{s=1}^i \bar{f}_j (X_1, \dots, \hat{t} \dots, X_{j+1}) (\sigma_1, \dots, [\lambda X_t, \sigma_s], \dots, \sigma_i) + \\
&+ \sum_{r < s} (-1)^{r+s} \bar{f}_j ([X_r, X_s], X_1, \dots, \hat{r} \dots \hat{s} \dots X_{j+1}) (\sigma_1, \dots, \sigma_i).
\end{aligned}$$

The cochain $\bar{f}_{j+1} (X_1, \dots, X_{j+1})$ is $C^\infty(M)$ -linear, i.e. belongs to the module $\Omega^{j+1}(M, \Lambda^{i-1} \mathbf{g}^* \otimes \xi)$. This gives

$$\begin{aligned}
& ((-1)^i d_{\nabla^+}^i \circ \Psi_{A,1}^{j,i} - \Psi_{A,1}^{j+1,i} \circ d_{A,1}^{j,i}) (f) (X_1, \dots, X_{j+1}) \\
&= (-1)^{(j+1)i} [d_{\nabla^+} ((-1)^j \bar{f}_{j+1} (X_1, \dots, X_{j+1}))] \\
&= 0.
\end{aligned}$$

If f has a compact support, the same hold for ρ_f and \bar{f}_{j+1} and we get the commutativity of the second diagram.

The next theorem is the main goal of this section. It describes the second terms of the spectral sequences (22) and (24) (see also [17]).

Theorem 5.10. *The homomorphisms*

$$\begin{aligned}
\Psi_{A,2} : E_{A,2}^{j,i} &\rightarrow H_{\nabla^+}^j(M, H_{\nabla^+}^i(\mathbf{g}, \xi)), \quad [f] \mapsto [(-1)^{ji} [\bar{f}_j]], \\
\Psi_{A_c,2} : E_{A_c,2}^{j,i} &\rightarrow H_{\nabla^+,c}^j(M, H_{\nabla^+}^i(\mathbf{g}, \xi)), \quad [f] \mapsto [(-1)^{ji} [\bar{f}_j]],
\end{aligned}$$

are isomorphisms of $C^\infty(M)$ -modules.

Proof. Clearly, we need to notice only that $\Psi_{A,2}$ is a composition of isomorphisms

$$\Psi_{A,2} : E_{A,2}^{j,i} \xrightarrow[\cong]{\sigma_{A,1}^{j,i}} H^j(E_{A,1}^{*,i}, d_{A,1}^{*,i}) \xrightarrow[\cong]{(\Psi_{A,1}^{j,i})^\#} H_{\nabla^+}^j(M, H_{\nabla^+}^i(\mathbf{g}, \xi))$$

and analogously for compact supports.

6 Algebroids and pairings

Assume that A is a transitive Lie algebroid with three representations

$$\nabla_r : A \rightarrow A(\xi_r), \quad r = 1, 2, 3,$$

and a pairing

$$F : \xi_1 \times \xi_2 \rightarrow \xi_3$$

compatible with the representations $(\nabla_1, \nabla_2, \nabla_3)$, i.e. fulfilling the property analogous to (5) in which we must replace X by $\gamma \in \Gamma(A)$. Then the multiplication of cochains

$$\wedge : \Lambda^j \mathbf{g}^* \otimes \xi_1 \times \Lambda^i \mathbf{g}^* \otimes \xi_2 \rightarrow \Lambda^{j+i} \mathbf{g}^* \otimes \xi_3$$

is compatible with

(a) suitable representations $\mathcal{L}_A^j, \mathcal{L}_A^i, \mathcal{L}_A^{j+i}$

$$\mathcal{L}_{A,\gamma}^{j+i}(f \wedge g) = \mathcal{L}_{A,\gamma}^{j+i}(f) \wedge g + f \wedge \mathcal{L}_{A,\gamma}^{j+i}(g),$$

$f \in \Gamma(\Lambda^j \mathbf{g}^* \otimes \xi_1), g \in \Gamma(\Lambda^i \mathbf{g}^* \otimes \xi_2)$,

(b) differentials $d_{\nabla_1^+}, d_{\nabla_2^+}, d_{\nabla_3^+}$

$$d_{\nabla_3^+}(f \wedge g) = d_{\nabla_1^+}(f) \wedge g + (-1)^j f \wedge d_{\nabla_2^+}(g),$$

f, g as above.

The latter equality gives the pairing of cohomology vector bundles

$$\wedge : H_{\nabla_1^+}^j(\mathbf{g}, \xi_1) \times H_{\nabla_2^+}^i(\mathbf{g}, \xi_2) \rightarrow H_{\nabla_3^+}^{j+i}(\mathbf{g}, \xi_3) \tag{28}$$

which is compatible with the suitable representations $\mathcal{L}_A^{\#,j}, \mathcal{L}_A^{\#,i}, \mathcal{L}_A^{\#,j+i}$ and finally with the flat covariant derivatives $\nabla^j, \nabla^i, \nabla^{j+i}$

$$\nabla_X^{j+i}([f] \wedge [g]) = \nabla_X^j([f]) \wedge [g] + [f] \wedge \nabla_X^i[g].$$

We assume in the sequel that $n = \text{rank } \mathbf{g}$ (and we recall that $m = \dim M$).

Together with three representations ∇_r one consider three graded filtered differential spaces $\Omega(A, \xi_1), \Omega_c(A, \xi_2), \Omega_c(A, \xi_3)$ (21), (23), and theirs spectral sequences $({}^1E_{A,s}^{j,i}, {}^1d_{A,\nabla,s}), ({}^2E_{A_c,s}^{j,i}, {}^2d_{A_c,\nabla,s}), ({}^3E_{A_c,s}^{j,i}, {}^3d_{A_c,\nabla,s})$. Using monomorphy of ρ_0 and of $\rho_{c,0}$, Lemma 5.1, we see immediately from the case of \mathbb{R} -linear cochains [9] that the following diagram commutes.

$$\begin{array}{ccc} {}^1E_{A,0}^{j,i} \times {}^2E_{A_c,0}^{j',i'} & \xrightarrow{\wedge} & {}^3E_{A_c,0}^{j+j',i+i'} \\ \downarrow a_A^{j,i} \times a_{A_c}^{j',i'} & & \downarrow a_A^{j,i} \times a_{A_c}^{j',i'} \\ \Omega^j(M, \Lambda^i \mathbf{g}^* \otimes \xi_1) \times \Omega_c^{j'}(\Lambda^{i'} \mathbf{g}^* \otimes \xi_2) & \xrightarrow{(-1)^{i'j} \cdot \wedge} & \Omega_c^{j+j'}(\Lambda^{i+i'} \mathbf{g}^* \otimes \xi_3). \end{array}$$

Passing twice to cohomology and using definitions of suitable homomorphisms we get the commutativity of the diagram

$$\begin{array}{ccc}
 {}^1E_{A,2}^{j,i} \times {}^2E_{A_c,2}^{j',i'} & \xrightarrow{\wedge} & {}^3E_{A_c,2}^{j+j',i+i'} \\
 \downarrow {}^1\Psi_{A,2}^{j,i} \times {}^2\Psi_{A_c,2}^{j',i'} & & \downarrow {}^3\Psi_{A_c,2}^{j+j',i+i'} \\
 H_{\nabla_1}^j(M, H_{\nabla_1^+}^i(\mathbf{g}, \xi_1)) \times H_{\nabla_{i',c}}^{j'}(M, H_{\nabla_2^+}^{i'}(\mathbf{g}, \xi_2)) & \xrightarrow{(-1)^{i'j} \cdot \wedge} & H_{\nabla_{i+i',c}}^{j+j'}(M, H_{\nabla_3^+}^{i+i'}(\mathbf{g}, \xi_3)).
 \end{array} \tag{29}$$

The main theorem of Chapter 1 one gets the very important **Conclusion 6.1.** *If ξ_3 is a line bundle and $(H_{\nabla_3^+}^n(\mathbf{g}, \xi_3), \nabla^n) \sim (or(M), \partial^{or})$ [in particular, $(\nabla_3^+)_x = \nabla_{\text{trad}} \otimes \text{id}$ according to Lemma 3.5] and the pairing of cohomology vector bundles*

$$\wedge : H_{\nabla_1^+}^i(\mathbf{g}, \xi_1) \times H_{\nabla_2^+}^{n-i}(\mathbf{g}, \xi_2) \rightarrow H_{\nabla_3^+}^n(\mathbf{g}, \xi_3)$$

is nondegenerate, then the same holds for the pairing

$$H_{\nabla_1}^j(M, H_{\nabla_1^+}^i(\mathbf{g}, \xi_1)) \times H_{\nabla_{n-i,c}}^{m-j}(M, H_{\nabla_2^+}^{n-i}(\mathbf{g}, \xi_2)) \rightarrow H_{\nabla_{n,c}}^m(M, H_{\nabla_3^+}^n(\mathbf{g}, \xi_3)) \xrightarrow{f_M^\#} \mathbb{R}. \tag{30}$$

i.e.

$$H_{\nabla_1}^j(M, H_{\nabla_1^+}^i(\mathbf{g}, \xi_1)) \cong (H_{\nabla_{n-i,c}}^{m-j}(M, H_{\nabla_2^+}^{n-i}(\mathbf{g}, \xi_2)))^*.$$

Diagram (29) assert that the nondegenerate pairing (30) is \pm equal to the multiplication of the second term of the spectral sequences

$${}^1E_{A,2}^{j,i} \times {}^2E_{A_c,2}^{m-j,n-i} \xrightarrow{\wedge} {}^3E_{A_c,2}^{m,n} \xrightarrow{\cong} \mathbb{R}$$

so the last is nondegenerate as well,

$${}^1E_{A,2}^{j,i} \cong ({}^2E_{A_c,2}^{m-j,n-i})^*,$$

and the main theorem of Chapter 4 gives that the multiplication of cohomology classes

$$\langle \cdot, \cdot \rangle_H : H_{\nabla_1}^j(A, \xi_1) \times H_{\nabla_{2,c}}^{m+n-j}(A, \xi_2) \rightarrow H_{\nabla_{3,c}}^{m+n}(A, \xi_3) \xrightarrow{\cong} \mathbb{R}$$

is nondegenerate too, i.e.

$$H_{\nabla_1}^j(A, \xi_1) \cong (H_{\nabla_{2,c}}^{m+n-j}(A, \xi_2))^*.$$

7 Evens-Lu-Weinstein pairing for transitive Lie algebroids

7.1 Nondegeneracy of Evens-Lu-Weinstein pairing for transitive Lie algebroids

We prove that for transitive Lie algebroid A the duality of Evens-Lu-Weinstein [4]

$$H^j(A) \times H_{D^{or},c}^{m+n-j}(A, Q_A^{or}) \rightarrow H_{D^{or},c}^{m+n}(A, Q_A^{or}) \rightarrow \mathbb{R}$$

is nondegenerate, i.e. $H_{D^{or},c}^{m+n}(A, Q_A^{or}) \xrightarrow{\cong} \mathbb{R}$ and

$$H^j(A) \cong (H_{D^{or},c}^{m+n-j}(A, Q_A^{or}))^*.$$

For arbitrary (nonregular in general) Lie algebroid A on a manifold M the authors [4] introduced a vector bundle

$$Q_A = \Lambda^{\text{top}} A \otimes \Lambda^{\text{top}} T^* M$$

(the notation Λ^{top} refers to the highest exterior power). Geometrically, sections of Q_A can be thought of as transverse measures to characteristic foliation $\text{Im } \#_A$ to any Lie algebroid A [4]. For Poisson manifolds, the Evens-Lu-Weinstein pairings takes the form of the pairing on the Poisson homology; for more applications see [4]. Ibidem, there is an example of nonregular Lie algebroid A over a compact oriented manifold for which the pairing $H^j(A) \times H_{D,c}^{m+n-j}(A, Q_A) \rightarrow \mathbb{R}$ is not necessarily nondegenerate. J.Huebschmann in [10] has generalized the construction of the bundle Q_A and the modular class θ_A to Lie-Rinehart algebras, an algebraic generalization of Lie algebroids.

We slightly modify the Weinstein construction to consider nonoriented manifolds:

$$Q_A^{or} = Q_A \otimes or(M).$$

For an oriented manifold M we can identify $Q_A^{or} = Q_A$.

In [4] a representation

$$D : A \rightarrow A(Q_A)$$

was introduced by

$$D_\gamma(Y \otimes \varphi) = L_\gamma(Y) \otimes \varphi + Y \otimes L_{\#_A(\gamma)}(\varphi),$$

$Y \in \Gamma(\Lambda^{\text{top}} A)$, $\varphi \in \Gamma(\Lambda^{\text{top}} T^* M) = \Omega^m(M)$, where $L_\gamma(Y) = [\gamma, Y]$ ($[\gamma, Y]$ denotes the Schouten bracket) and $L_{\#_A(\gamma)}(\varphi)$ is the usual Lie derivative of a differential form φ . We recall that for $Y = \gamma_1 \wedge \dots \wedge \gamma_t$

$$L_\gamma(Y) = \sum_i \gamma_1 \wedge \dots \wedge [[\gamma, \gamma_i]] \wedge \dots \wedge \gamma_t.$$

There is some interest to consider the representation D in the context of intrinsic characteristic classes of Lie algebroids [3], [5].

We modify the representation D to

$$D^{or} = D \otimes \partial_A^{or} : A \rightarrow A(Q_A^{or}).$$

In the sequel we will be interested only in the transitive case. A choice of a connection $\lambda : TM \rightarrow A$ enables us to identify

$$\begin{aligned} \Lambda^{m+n} A &= \Lambda^n \mathbf{g} \otimes \Lambda^m TM, \\ \varepsilon \wedge (\Lambda^m \lambda)(X) &= \varepsilon \otimes X, \end{aligned}$$

and

$$Q_A^{or} = \Lambda^n \mathfrak{g} \otimes \Lambda^m TM \otimes \Lambda^m T^*M \otimes or(M) = \Lambda^n \mathfrak{g} \otimes or(M). \tag{31}$$

Lemma 7.1. (a) $D^+ : \mathfrak{g} \rightarrow \text{End}(Q_A)$ is defined by

$$D_\sigma^+ = (\nabla_{\text{trad}})_\sigma = \text{tr}(\text{ad}_\sigma) \cdot \text{id}, \quad \sigma \in \Gamma(\mathfrak{g}).$$

(b) $H_{D^+}^n(\mathfrak{g}, Q_A) = \Lambda^n \mathfrak{g}^* \otimes Q_A, H_{D^{or}}^n(\mathfrak{g}, Q_A^{or}) = \Lambda^n \mathfrak{g}^* \otimes Q_A^{or}.$

Proof. (a) Consider locally defined nonsingular section of Q_A of the form $\varepsilon_U \otimes X_U \otimes \varphi_U$, $\varepsilon_U \in \Gamma(\Lambda^n \mathfrak{g}_U), X_U \in \Gamma(\Lambda^m TM_U), \varphi_U \in \Gamma(\Lambda^m T^*M_U)$, and assume that $\langle X_U, \varphi_U \rangle = 1$. For $\sigma \in \Gamma(\mathfrak{g})$ we have $\#_A(\sigma) = 0$ and $[[\sigma, \lambda W_i]] \in \Gamma(\mathfrak{g})$. Therefore if $\varepsilon_U = \sigma_1 \wedge \dots \wedge \sigma_n, X_U = W_1 \wedge \dots \wedge W_m, \sigma_i \in \Gamma(\mathfrak{g}), W_i \in \Gamma(TM_U)$,

$$\begin{aligned} D_\sigma(\varepsilon_U \otimes X_U \otimes \varphi_U) &= L_\sigma(\varepsilon_U \otimes X_U) \otimes \varphi_U \\ &= L_\sigma(\sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m) \otimes \varphi_U \\ &= \sum_i \sigma_1 \wedge \dots \wedge [[\sigma, \sigma_i]] \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes \varphi_U \\ &= \text{tr}(\text{ad}_\sigma) \cdot \varepsilon_U \otimes X_U \otimes \varphi_U. \end{aligned}$$

(b) Follows immediately from Proposition 3.2.

The vector bundle $\Lambda^n \mathfrak{g}^* \otimes Q_A$ is trivial. Indeed, the classical homomorphism

$$\begin{aligned} c : \Lambda^n \mathfrak{g}^* \otimes Q_A &= \Lambda^n \mathfrak{g}^* \otimes \Lambda^n \mathfrak{g} \otimes \Lambda^m TM \otimes \Lambda^m T^*M \rightarrow M \times \mathbb{R} \\ c(\varepsilon^* \otimes \varepsilon \otimes X \otimes \varphi) &= \langle \varepsilon^*, \varepsilon \rangle \cdot \langle X, \varphi \rangle \end{aligned} \tag{32}$$

is an isomorphism. Therefore

$$c \otimes \text{id} : \Lambda^n \mathfrak{g}^* \otimes Q_A^{or} \xrightarrow{\cong} or(M). \tag{33}$$

Let $A(c) : A(\Lambda^n \mathfrak{g}^* \otimes Q_A) \rightarrow A(M \times \mathbb{R})$ be the induced isomorphism of Lie algebroids [12],

$$A(c)(u)(f) = c(u(c^{-1} \circ f)),$$

$u \in \Gamma(A(\Lambda^n \mathfrak{g}^* \otimes Q_A)), f \in C^\infty(M)$. Let

$$\nabla^D : TM \rightarrow A(H_{D^+}^n(\mathfrak{g}, Q_A)) = A(\Lambda^n \mathfrak{g}^* \otimes Q_A)$$

be the induced flat adjoint covariant derivative (27) for D . Analogously we have $\nabla^{D^{or}} : TM \rightarrow A(\Lambda^n \mathfrak{g}^* \otimes Q_A^{or})$.

Lemma 7.2. *The compositions*

$$\begin{aligned} \nabla^D : TM &\rightarrow A(\Lambda^n \mathfrak{g}^* \otimes Q_A) \xrightarrow[\cong]{A(c)} A(TM \times \mathbb{R}), \\ \nabla^{D^{or}} : TM &\rightarrow A(\Lambda^n \mathfrak{g}^* \otimes Q_A^{or}) \xrightarrow[\cong]{A(c \otimes \text{id})} A(or(M)), \end{aligned}$$

are trivial representations ∂ and ∂^{or} , respectively, so

$$\begin{aligned} (H_{D^+}^n(\mathbf{g}, Q_A), \nabla^D) &\sim (TM \times \mathbb{R}, \partial), \\ (H_{D^{or+}}^n(\mathbf{g}, Q_A), \nabla^{D^{or}}) &\sim (or(M), \partial^{or}). \end{aligned}$$

Proof. It is necessary to check it locally. Take locally defined nonsingular sections $\varepsilon_U \in \Gamma(\Lambda^n \mathbf{g}_U)$, $X_U \in \Gamma(\Lambda^m T M_U)$ and their duals $\varepsilon_U^* \in \Gamma(\Lambda^n \mathbf{g}_U^*)$, $\varphi_U \in \Gamma(\Lambda^m T^* M_U)$, $\langle \varepsilon_U^*, \varepsilon_U \rangle = 1$, $\langle X_U, \varphi_U \rangle = 1$. On the set U arbitrary section of the bundle $\Lambda^n \mathbf{g}^* \otimes Q_A = \Lambda^n \mathbf{g}^* \otimes \Lambda^n \mathbf{g} \otimes \Lambda^m T M \otimes \Lambda^m T^* M$ is of the form $f \cdot \varepsilon_U^* \otimes \varepsilon_U \otimes X_U \otimes \varphi_U$, $f \in C^\infty(U)$. For $X \in \mathfrak{X}(M_U)$, $X_U = W_1 \wedge \dots \wedge W_m$ (W_1, \dots, W_m is a base of vector fields on U) and $\varphi_U = W_1^* \wedge \dots \wedge W_m^*$, W_i^* is the dual basis, and $\varepsilon_U = \sigma_1 \wedge \dots \wedge \sigma_n$ (σ_i is a base of the vector bundle \mathbf{g} on U), we write $[[\lambda X, \sigma_i]] = \sum_j g_i^j \cdot \sigma_j$, $[[\lambda X, \lambda W_i]] = \sum_k h_i^k \cdot \sigma_k + a_i^k \cdot \lambda_k$, so $[X, W_i] = a_i^k \cdot W_k$. Then

$$\begin{aligned} D_{\lambda X}(\varepsilon_U \otimes X_U \otimes \varphi_U) &= L_{\lambda X}(\varepsilon_U \otimes X_U) \otimes \varphi_U + \varepsilon_U \otimes X_U \otimes L_X \varphi_U \\ &= \left(\sum_i g_i^i \cdot \sigma \wedge \lambda X \right) \otimes \varphi_U + \sigma \wedge \left(\sum_i a_i^i \cdot \lambda X \right) \otimes \varphi_U + \sigma \wedge \lambda X \otimes \sum (-a_i^i) \varphi_U \\ &= \sum_i g_i^i \cdot \varepsilon_U \otimes X_U \otimes \varphi_U. \end{aligned}$$

Therefore

$$\begin{aligned} \nabla_X^D(f \cdot \varepsilon_U^* \otimes \varepsilon_U \otimes X_U \otimes \varphi_U)(\varepsilon_U) &= D_{\lambda X}(f \cdot \varepsilon_U \otimes X_U \otimes \varphi_U) - \\ &\quad - \sum_i f \cdot \varepsilon_U^* (\sigma_1 \wedge \dots \wedge [[\lambda X, \sigma_i]] \wedge \dots \wedge \sigma_n) \cdot \varepsilon_U \otimes X_U \otimes \varphi_U \\ &= \partial_X f \cdot \varepsilon_U \otimes X_U \otimes \varphi_U + f \cdot D_{\lambda X}(\varepsilon_U \otimes X_U \otimes \varphi_U) - f \cdot \sum_i g_i^i \cdot \varepsilon_U \otimes X_U \otimes \varphi_U \\ &= \partial_X f \cdot \varepsilon_U \otimes X_U \otimes \varphi_U \\ &= \partial_X f \cdot (\varepsilon_U^* \otimes \varepsilon_U \otimes X_U \otimes \varphi_U)(\varepsilon_U). \end{aligned}$$

Finally

$$\begin{aligned} (A(c) \circ \nabla_X^D)(f) &= c(\nabla_X^D(f \cdot \varepsilon_U^* \otimes \varepsilon_U \otimes X_U \otimes \varphi_U)) \\ &= c(\partial_X f \cdot (\varepsilon_U^* \otimes \varepsilon_U \otimes X_U \otimes \varphi_U)) \\ &= \partial_X f. \end{aligned}$$

For the proof of the second part we notice that for local ∂^{or} -constant section σ_0 of $or(M_U)$ one has

$$\begin{aligned} (A(c \otimes id) \circ \nabla_X^{D^{or}})(f \otimes \sigma_0) &= (A(c) \circ \nabla_X^D)(f) \otimes \sigma_0 \\ &= \partial_X f \otimes \sigma_0 \\ &= \partial_X^{or}(f \otimes \sigma_0). \end{aligned}$$

Theorem 7.3. *For an arbitrary transitive Lie algebroid A*

$$H_{D^{or},c}^{m+n}(A, Q_A^{or}) \cong \mathbb{R},$$

and the Evens-Lu-Weinstein cohomology pairing

$$H^j(A) \times H_{D^{or},c}^{m+n-j}(A, Q_A^{or}) \rightarrow H_{D^{or},c}^{m+n}(A, Q_A^{or}) \cong \mathbb{R}$$

is nondegenerate, i.e.

$$H^j(A) \cong (H_{D^{or},c}^{m+n-j}(A, Q_A^{or}))^*.$$

Proof. Theorem 3.4 and Lemma 7.1 show that the pairing

$$H^i(\mathfrak{g}) \times H_{D^{or+}}^{n-i}(\mathfrak{g}, Q_A^{or}) \rightarrow H_{D^{or+}}^n(\mathfrak{g}, Q_A^{or})$$

is nondegenerate. On account of Theorem 2.3 and Conclusion 6.1 we assert that the pairing

$$H_{\nabla^D}^j(M, H^i(\mathfrak{g})) \times H_{\nabla^{or},c}^{m-j}(M, H_{D^{or+}}^{n-i}(\mathfrak{g}, Q_A^{or})) \rightarrow H_{\nabla^{D^{or},c}}^m(M, H_{D^{or+}}^n(\mathfrak{g}, Q_A^{or})) \xrightarrow{\cong} \mathbb{R},$$

is nondegenerate. Equivalently, this is a multiplication of the second terms of the Hochschild-Serre spectral sequences of graded filtered differential spaces $\Omega(A)$ with the trivial differential and $\Omega_c(A, Q_A^{or})$ with the differential D^{or} . The fundamental Theorem 4.4, see also mentioned above Conclusion 6.1, completes the proof.

7.2 Remarks on the top group of cohomology

Analyzing the proof of Theorem 4.4 and composing isomorphism (33) with isomorphism (4) we can define the isomorphism $I : H_{D^{or},c}^{m+n}(A, Q_A^{or}) \xrightarrow{\cong} \mathbb{R}$ as a composition

$$\begin{aligned} I : H_{D^{or},c}^{m+n}(A, Q_A^{or}) &= H_{D^{or},c}^{m,n}(A, Q_A^{or}) = E_0^{m,n}(H_{D^{or},c}(A, Q_A^{or})) \xleftarrow[\cong]{\sigma_{A_c,\infty}^{m,n}} E_{A_c,\infty}^{m,n} = \\ &= E_{A_c,2}^{m,n} \xrightarrow[\cong]{\Psi_{A_c,2}^{m,n}} H_{\nabla^{D^{or},c}}^m(M, H_{D^{or+}}^n(\mathfrak{g}, Q_A^{or})) = \\ &= H_{\nabla^{D^{or},c}}^m(M, \Lambda^n \mathfrak{g}^* \otimes Q_A^{or}) \xrightarrow[\cong]{(c \otimes id)^\#} H_{\partial^{or},c}^m(M, or(M)) \xrightarrow[\cong]{f_M^{or,\#}} \mathbb{R}. \end{aligned}$$

We compare this isomorphism with the one defined by direct formula in [E-L-W] restricting our interest to transitive Lie algebroids. Immediately from the definition of $\Psi_{A_c,2}^{m,n}$ (see Theorem 5.10), $\Psi_{A_c,2}^{m,n}[f] = (-1)^{mn} [\bar{f}_m]$, and definition of $\sigma_{A_c,\infty}^{m,n}$ we observe that

$$I_1 : H_{D^{or},c}^{m+n}(A, Q_A^{or}) = E_0^{m,n} \xleftarrow[\cong]{\sigma_{A_c,\infty}^{m,n}} E_{A_c,\infty}^{m,n} = E_{A_c,2}^{m,n} \xrightarrow[\cong]{\Psi_{A_c,2}^{m,n}} H_{\nabla^{D^{or},c}}^m(M, \Lambda^n \mathfrak{g}^* \otimes Q_A^{or})$$

is given by the formula looking analogously to $\Psi_{A_c,2}^{m,n}$, $I_1[f] = (-1)^{mn} [\bar{f}_m]$, or equivalently (under the identification $\Lambda^{m+n} A^* = \Lambda^m T^*M \otimes \Lambda^n \mathfrak{g}^*$ given by the help of a connection $\lambda : TM \rightarrow A$) by

$$I_1((\varphi \otimes \varepsilon^*) \otimes q) = (-1)^{mn} [\varphi \otimes (\varepsilon^* \otimes q)]$$

where $\varphi \in \Omega_c^m(M)$, $\varepsilon^* \in \Gamma(\Lambda^n \mathfrak{g}^*)$, $q \in \Gamma(Q_A^{or})$. Therefore if $q = \varepsilon \otimes X \otimes \mu \otimes e$, $\varepsilon \in \Gamma(\Lambda^n \mathfrak{g})$, $X \in \Gamma(\Lambda^m TM)$, $\mu \in \Gamma(\Lambda^m T^*M) = \Omega^m(M)$, $e \in \Gamma(or(M))$ then

$$(c \otimes id)_{\#} \circ I_1((\varphi \otimes \varepsilon^*) \otimes \varepsilon \otimes X \otimes \mu \otimes e) = (-1)^{mn} [\varphi \cdot \langle \varepsilon^*, \varepsilon \rangle \cdot \langle X, \mu \rangle \otimes e].$$

So, for $f = (\varphi \otimes \varepsilon^*) \otimes \varepsilon \otimes X \otimes \mu \otimes e$ we get

$$\begin{aligned} I[f] &= (-1)^{mn} \int_M^{or} \langle \varepsilon^*, \varepsilon \rangle \cdot \langle X, \mu \rangle \cdot \varphi \otimes e = (-1)^{mn} \int_M^{or} \langle \varepsilon^*, \varepsilon \rangle \cdot \langle X, \varphi \rangle \cdot \mu \otimes e \\ &= (-1)^{mn} \int_M^{or} \langle \varphi \otimes \varepsilon^*, X \otimes \varepsilon \rangle \cdot \mu \otimes e \end{aligned}$$

which is concordant up to the sign with the definition of Evens-Lu-Weinstein [4] given by them only for oriented compact manifold (but for any Lie algebroid, not necessary transitive).

The fact $H_{D^{or,c}}^{m+n}(A, Q_A) \xrightarrow{\cong} \mathbb{R}$ for transitive Lie algebroids is not proved in [4]. Below we prove this immediately without use of the spectral sequences.

(a) *on oriented manifolds.* The authors of [4] introduced an isomorphism of vector bundles

$$\begin{aligned} \tilde{\rho} : \Lambda^{\text{top}} A^* \otimes \Lambda^{\text{top}} A \otimes \Lambda^{\text{top}} T^*M &\rightarrow \Lambda^{\text{top}} T^*M, \\ \tilde{\rho}(\Psi \otimes Y \otimes \mu) &= \langle \Psi, Y \rangle \cdot \mu \end{aligned}$$

and proved a version of Stokes Theorem (to be sure for compact manifold but without troubles we can extend it to differential forms with compact support on arbitrary oriented manifold).

Theorem 7.4 (Stokes Theorem [4]). *Let $\text{rank } A = r$. For $r - 1$ -form $\Psi' \in \Gamma(\Lambda^{r-1} A^*)$ we have*

$$\tilde{\rho}(d_D(\Psi' \otimes Y \otimes \mu)) = (-1)^{r-1} d_{dR}(\iota_{\#A}(\Psi' \lrcorner Y)\mu).$$

Consequently, if the form $\Psi' \otimes Y \otimes \mu$ has compact support then

$$\int_M \tilde{\rho}(d_D(\Psi' \otimes Y \otimes \mu)) = 0.$$

Put

$$\tilde{\rho}^{r-1} : \Lambda^{r-1} A^* \otimes Q_A \rightarrow \Lambda^{m-1} T^*M, \quad (\Psi' \otimes Y \otimes \mu) \mapsto (-1)^{r-1} \iota_{\#A}(\Psi' \lrcorner Y)\mu,$$

and notice the commutativity of the diagram

$$\begin{array}{ccc} \Omega_c^{r-1}(A, Q_A) & \xrightarrow{\tilde{\rho}_c^{r-1}} & \Omega_c^{m-1}(M) \\ \downarrow d_D & & \downarrow d_{dR} \\ \Omega_c^r(A, Q_A) & \xrightarrow{\tilde{\rho}_c} & \Omega_c^m(M). \end{array}$$

From this we deduce that $\tilde{\rho}_c$ induces an \mathbb{R} -linear homomorphism in cohomology

$$\tilde{\rho}_{c,\#} : H_{D,c}^r(A, Q_A) \rightarrow H_c^m(M).$$

Since $\tilde{\rho}$ is an isomorphism $\tilde{\rho}_{c,\#}$ is an epimorphism.

Lemma 7.5. *If A is transitive Lie algebroid, then $\tilde{\rho}_{c,\#}$ is an isomorphism.*

Proof. One can easily see the lemma provided that $\tilde{\rho}^{r-1}$ is an epimorphism. It is a simple matter to show that $\tilde{\rho}_x^{r-1}$ is an epimorphism at every point $x \in M$ using transitivity of the Lie algebroid A i.e. using the fact that the anchor $(\#_A)_x : A_x \rightarrow T_x M$ is an epimorphism. This finishes the proof that

$$\tilde{\rho}_{c,\#} : H_{D,c}^r(A, Q_A) \rightarrow H_c^m(M) \xrightarrow[\cong]{\int_M^\#} \mathbb{R}$$

is an isomorphism.

(b) *on nonoriented manifolds.* We prove this analogously multiplying the vector bundles by $or(M)$ and use the Stokes theorem for densities [1].

7.3 Exceptional property of the Evens-Lu-Weinstein representation

Assume A is a transitive Lie algebroid. Before the next theorems we must give algebroid's equivalent of some lemmas from Chapter 1. For any A -connection $\nabla : A \rightarrow A(\xi)$ and a 1-form $\omega \in \Omega^1(A)$ we define a new A -connection

$$\nabla^\omega : A \rightarrow A(\xi), \quad \nabla_\gamma^\omega \nu = \nabla_\gamma \nu + \omega(\gamma) \cdot \nu.$$

The curvature tensors $R^{\nabla^\omega}, R^\nabla \in \Omega^2(A, \xi)$ of the connections ∇^ω and ∇ are related via the formula

$$R^{\nabla^\omega} = R^\nabla + d_A \omega \otimes id.$$

Therefore, if ∇ is flat (it means, ∇ is a representation) then ∇^ω is flat if and only if ω is closed. Each A -connection $\nabla : A \rightarrow A(M \times \mathbb{R})$ in the trivial vector bundle $M \times \mathbb{R}$ is of the form ∂_A^ω , indeed, we need to put $\omega(\gamma) = \nabla_\gamma(1)$.

For a line bundle ξ and a representation $\nabla : A \rightarrow A(\xi)$ the differential equation $\nabla \nu = 0$ is locally uniquely integrable provided that it is locally integrable.

Lemma 7.6. *For a line bundle ξ and a representation $\nabla : A \rightarrow A(\xi)$ the differential equation $\nabla \nu = 0$ is locally integrable if and only if $\nabla^+ = 0$. This last condition is equivalent to the projectability of ∇ , i.e. that $\nabla = \tilde{\nabla} \circ \#_A$ for some usual flat covariant derivative $\tilde{\nabla}$ on M in the vector bundle ξ .*

Proof. "⇒" Assume that $\nabla \nu = 0$ is locally integrable. If ν is locally defined nonsingular ∇ -constant section of ξ then arbitrary section is equal to $\nu_1 = f \cdot \nu$ for a smooth function f and for $\sigma \in \Gamma(\mathfrak{g})$

$$\nabla_\sigma^+(f \cdot \nu) = \partial_{\#_A(\sigma)} f \cdot \nu + f \cdot \nabla_\sigma \nu = 0.$$

"⇐" Assume that $\nabla^+ = 0$. Take $x_0 \in M$ and $u \in \xi_{x_0}$. Locally $(x_0 \in U \cong \mathbb{R}^m)$ $\nabla_U = \partial_{A_U}^\omega : A_U \rightarrow A(\xi_U) = A(U \times \mathbb{R})$ for a closed 1-form $\omega \in \Omega^1(A_U)$. By assumption

$\nabla^+ = 0$, $\omega(\sigma) \cdot \nu = 0$ for all $\sigma \in \Gamma(\mathfrak{g})$ and $\nu \in \Gamma(\xi)$, so $\omega(\sigma) = 0$ and ω is projectable on U , $\omega = \#_A^*(\bar{\omega})$ for some $\bar{\omega} \in \Omega^1(U)$. Since the anchor $\#_A$ is an epimorphism, the pullback of the differential forms $\#_A^*$ is a monomorphism. Therefore, since $0 = d_A\omega = d_A(\#_A^*(\bar{\omega})) = \#_A^*(d_{dR}\bar{\omega})$ we get $d_{dR}\bar{\omega} = 0$. Clearly, then $\bar{\omega} = df$ for some function $f \in C^\infty(U)$. It is easy to see that the section $\sigma = e^{-f}$ of the bundle $U \times \mathbb{R} \cong \xi_U$ is ∇_U -constant.

Similar considerations show that for trivial vector bundle $\xi = M \times \mathbb{R}$ and a representation ∂_A^ω the following conditions are equivalent (1) $\nabla^+ = 0$, (2) ω is projectable (i.e. $\omega = \#_A^*(\bar{\omega})$ for some $\bar{\omega} \in \Omega^1(M)$). On the other hand, if ω is exact, i.e. $0 = [\omega]_A \in H^1(A)$, then $\nabla^+ = 0$, which implies that the differential equation $\nabla\nu = 0$ is locally uniquely integrable.

By the definition the 0-group of cohomology can be written similarly to (1).

$$H_\nabla^0(A, \xi) = \{\nu \in \Gamma(\xi); \nabla\nu = 0\}.$$

Proposition 7.7. (1) $H_\nabla^0(A, \xi) = 0$ if ξ is nontrivial.

(2) For the trivial vector bundle $\xi = M \times \mathbb{R}$ and $\nabla = \partial_A^\omega$ for closed 1-form $\omega \in \Omega^1(A)$ we have

$$H_\nabla^0(A, \xi) \neq 0 \iff [\omega]_A = 0.$$

In particular, if $H_\nabla^0(A, \xi) \neq 0$ then $\nabla^+ = 0$.

Proof. (1) Evidently, since each section of nontrivial line bundle ξ is singular and by Lemma 7.6 the set $\{x; \nabla\nu = 0\}$ is open-closed.

(2) This result may be proved in the same way as in the case of $A = TM$, i.e. as the formula (3), see also Example 2.6 from Chapter 1.

Proposition 7.7(1) generalizes observation (●●) from section 2.1.

Proposition 7.8. Let ξ be a line bundle and fix an isomorphism $\varphi : \xi \otimes \xi \rightarrow M \times \mathbb{R}$. Let us assume that φ transforms a given A -representation $\nabla : A \rightarrow A(\xi)$ to A -representation ∂_A^ω for a closed 1-form $\omega \in \Omega^1(A)$. Then there exists a linear isomorphism

$$H_{D^{or} \otimes \nabla - \omega}^j(A, Q_A^{or} \otimes \xi) \cong H_{\nabla, c}^{m+n-j}(A, \xi)^*.$$

In particular

$$H_{D^{or} \otimes \nabla - \omega}^0(A, Q_A^{or} \otimes \xi) \cong H_{\nabla, c}^{m+n}(A, \xi)^*.$$

Proof. Consider the multiplication by reals

$$\rho : Q_A^{or} \times (M \times \mathbb{R}) \rightarrow Q_A^{or}.$$

ρ is compatible with $(D^{or}, \partial_A^\omega, (D^{or})^\omega)$. The canonical nondegenerate pairing

$$F : Q_A^{or} \otimes \xi \times \xi \rightarrow Q_A^{or} \otimes \xi \otimes \xi$$

is compatible with $(D^{or} \otimes \nabla, \nabla, D^{or} \otimes \nabla \otimes \nabla)$, so the composition

$$\tilde{F} : Q_A^{or} \otimes \xi \times \xi \xrightarrow{F} Q_A^{or} \otimes \xi \otimes \xi \xrightarrow{id \otimes \varphi} Q_A^{or} \otimes (M \times \mathbb{R}) \xrightarrow{\rho} Q_A^{or}$$

is compatible with $(D^{or} \otimes \nabla, \nabla, (D^{or})^\omega)$ which implies that it is also compatible with $(D^{or} \otimes \nabla^{-\omega}, \nabla, D^{or})$. Therefore, for each point $x \in M$, the pairing $\tilde{F}_x : Q_{A,x}^{or} \otimes \xi_x \times \xi_x \rightarrow Q_{A,x}^{or}$ is compatible with the representations $((D^{or} \otimes \nabla^{-\omega})_x^+, \nabla_x^+, D_x^{or+})$ of the isotropy Lie algebra \mathfrak{g}_x in the vector spaces $Q_{A,x}^{or} \otimes \xi_x$, ξ_x , $Q_{A,x}^{or}$, respectively. From this it follows that the differentials $d_{(D^{or} \otimes \nabla^{-\omega})_x^+}$, $d_{\nabla_x^+}$, $d_{(D^{or})_x^+}$ fulfil condition (6) from Lemma 3.1. Of course, $d_{(D^{or})_x^+} = d_{\text{trad}} \otimes id$ satisfies condition (3) from the mentioned lemma. Since

$$\wedge : \Lambda^i \mathfrak{g}_x^* \otimes (Q_{A,x}^{or} \otimes \xi_x) \times \Lambda^{n-i} \mathfrak{g}_x^* \otimes \xi_x \rightarrow \Lambda^n \mathfrak{g}_x^* \otimes Q_{A,x}^{or}$$

is nondegenerate, the generalized Chern-Hirzebruch-Serre Lemma 3.1 asserts that induced pairing in cohomology

$$H_{(D^{or} \otimes \nabla^{-\omega})^+}^i(\mathfrak{g}, Q_A^{or} \otimes \xi) \times H_{\nabla^+}^{n-i}(\mathfrak{g}, \xi) \rightarrow H_{D^{or+}}^n(\mathfrak{g}, Q_A^{or}) \xrightarrow{1.(7.2)} or(M)$$

is nondegenerate at every point $x \in M$. The fundamental Theorem 4.4, see also Conclusion 6.1, shows that the pairing

$$H_{D^{or} \otimes \nabla^{-\omega}}^j(A, Q_A^{or} \otimes \xi) \times H_{\nabla, c}^{m+n-j}(A, \xi) \rightarrow H_{D^{or}, c}^{m+n}(A, Q_A^{or})$$

is nondegenerate. This ends the proof.

Conclusion 7.9. *If ξ is not isomorphic to Q_A^{or} (i.e. ξ is not isomorphic to $\Lambda^n \mathfrak{g} \otimes or(M)$), see (31), then for an arbitrary connection $\nabla : A \rightarrow A(\xi)$ we have $H_{\nabla, c}^{m+n}(A, \xi) = 0$.*

Proof. If ξ is not isomorphic to Q_A^{or} the vector bundle $Q_A^{or} \otimes \xi$ is not trivial so Proposition 7.7 gives $H_{D^{or} \otimes \nabla^{-\omega}}^0(A, Q_A^{or} \otimes \xi) = 0$. Proposition 7.8 proves our theorem.

The next theorem is one of the importest theorems of the paper. Compare this theorem and Theorem 7.3 with Theorem 5.4 from [10].

Theorem 7.10. *For a line bundle ξ and a representation $\nabla : A \rightarrow A(\xi)$ the following conditions are equivalent:*

- (a) $H_{\nabla, c}^{m+n}(A, \xi) \neq 0$,
- (b) $H_{\nabla, c}^{m+n}(A, \xi) = \mathbb{R}$ and the pairing $H^j(A) \times H_{\nabla, c}^{m+n-j}(A, \xi) \rightarrow H_{\nabla, c}^{m+n}(A, \xi) \cong \mathbb{R}$ is nondegenerate, i.e. $H^j(A) \cong (H_{\nabla, c}^{m+n-j}(A, \xi))^*$,
- (c) $(\xi, \nabla) \sim (Q_A^{or}, D^{or})$.

Proof. (c) \implies (b) by Lemma 7.2, (b) \implies (a) is evident.

(a) \implies (c). Let $H_{\nabla, c}^{m+n}(A, \xi) \neq 0$. By Conclusion 7.9 $\xi \cong Q_A^{or}$. It remains to compare the representations. Consider then a flat bundle $(\xi = Q_A^{or}, \nabla)$ and any linear isomorphism

$$Q_A^{or} \otimes Q_A^{or} \xrightarrow{\cong} M \times \mathbb{R}, \quad \nabla \otimes \nabla \sim \partial_A^\omega.$$

By Proposition 7.8

$$H_{D^{or} \otimes \nabla^{-\omega}}^0(A, Q_A^{or} \otimes Q_A^{or}) \cong H_{\nabla, c}^{m+n}(A, Q_A^{or}) \neq 0,$$

therefore there exists a nonsingular global section $\nu \in \Gamma(Q_A^{or} \otimes Q_A^{or})$ which is $D^{or} \otimes \nabla^{-\omega}$ -constant. Additionally, $\nabla \otimes \nabla \sim \partial_A^\omega$ implies $\nabla \otimes \nabla^{-\omega} \sim \partial_A$ which means that there exists a second nonsingular section $\nu' \in \Gamma(Q_A^{or} \otimes Q_A^{or})$ which is $\nabla \otimes \nabla^{-\omega}$ -constant. The bundle $Q_A^{or} \otimes Q_A^{or}$ is 1-dimensional, so $\nu' = f \cdot \nu$ for a nonsingular function $f \in C^\infty(M)$. Write locally $\nu = \nu'_\alpha \otimes \nu_\alpha$ for nonsingular sections ν'_α, ν_α of Q_A^{or} . Then

$$\begin{aligned} 0 &= (D^{or} \otimes \nabla^{-\omega})_\gamma(\nu) = D_\gamma^{or}(\nu'_\alpha) \otimes \nu_\alpha + \nu'_\alpha \otimes \nabla_\gamma^{-\omega}(\nu_\alpha), \\ 0 &= (\nabla \otimes \nabla^{-\omega})_\gamma(f \cdot \nu) = \nabla_\gamma(f \cdot \nu'_\alpha) \otimes \nu_\alpha + f \cdot \nu'_\alpha \otimes \nabla_\gamma^{-\omega}(\nu_\alpha). \end{aligned}$$

Multiplying first equation by f and then subtracting the second we get

$$(f \cdot D_\gamma^{or}(\nu'_\alpha) - \nabla_\gamma(f \cdot \nu'_\alpha)) \otimes \nu_\alpha = 0.$$

The nonsingularity of ν_α yields the equation $f \cdot D_\gamma^{or}(\nu'_\alpha) = \nabla_\gamma(f \cdot \nu'_\alpha)$. The bundle Q_A^{or} is 1-dimensional, so

$$f \cdot D_\gamma^{or}(\bar{\nu}) = \nabla_\gamma(f \cdot \bar{\nu}) \tag{34}$$

for all $\bar{\nu} \in \Gamma(Q_A^{or})$. Define a linear isomorphism

$$\varphi : Q_A^{or} \rightarrow Q_A^{or}, \quad \bar{\nu} \mapsto f \cdot \bar{\nu}.$$

By (34) one has that $(Q_A^{or}, \nabla) \sim (Q_A^{or}, D^{or})$.

7.4 Characterization of transitive Lie algebroids with Poincaré duality

The last aim is to characterize two classes of transitive Lie algebroids.

- (1) $H_{\partial_A, c}^{m+n}(A) \neq 0$ - the top group of real compact cohomology is not trivial.

This condition is equivalent to $(Q_A^{or}, D^{or}) \sim (M \times \mathbb{R}, \partial_A)$. These classes fulfil the Poincaré duality: the pairing

$$H^j(A) \times H_c^{m+n-j}(A) \rightarrow H_c^{m+n}(A) \cong \mathbb{R}$$

is not degenerate, see Theorem 7.3, i.e. $H^j(A) \cong (H_c^{m+n-j}(A))^*$.

- (2) $H_{\partial_A^{or}, c}^{m+n}(A, or(M)) \neq 0$ - the top group of $or(M)$ -valued compact cohomology is not trivial.

This condition is equivalent to $(Q_A^{or}, D^{or}) \sim (or(M), \partial_A^{or})$. In this class the multiplication of cohomology classes

$$H^j(A) \times H_{\partial_A^{or}, c}^{m+n-j}(A, or(M)) \rightarrow H_{\partial_A^{or}, c}^{m+n}(A, or(M)) \cong \mathbb{R}$$

is not degenerate, see Theorem 7.3, i.e. $H^j(A) \cong (H_{\partial_A^{or}, c}^{m+n-j}(A))^*$.

Before the characterization of these classes we reduce the Evens-Lu-Weinstein representation (Q_A^{or}, D^{or}) to equivalent simple form (only for transitive Lie algebroids of course). We recall that the adjoint representation $ad_A : A \rightarrow A(\mathfrak{g})$ induces a top-power representation $ad_A^{top} : A \rightarrow A(\Lambda^{top} \mathfrak{g})$ by

$$(ad_A^{top})_\gamma(\sigma_1 \wedge \dots \wedge \sigma_n) = \sum_i \sigma_1 \wedge \dots \wedge [\gamma, \sigma_i] \wedge \dots \wedge \sigma_n = \sum_i a_i^i \cdot \sigma_1 \wedge \dots \wedge \sigma_n$$

where $[\gamma, \sigma_i] = \sum_j a_i^j \cdot \sigma_j$.

Lemma 7.11. *There exist isomorphisms of flat vector bundles*

$$\begin{aligned} (Q_A, D) &\cong (\Lambda^{top} \mathfrak{g}, ad_A^{top}), \\ (Q_A^{or}, D^{or}) &\cong (\Lambda^{top} \mathfrak{g} \otimes or(M), ad_A^{top} \otimes \partial_A^{or}). \end{aligned}$$

Proof. It is necessary to show the first assertion, because the second follows from first by tensor product with $or(M)$. Fix arbitrary a connection $\lambda : TM \rightarrow A$ and a linear isomorphism

$$\begin{aligned} \bar{\mathcal{K}} : \Lambda^{m+n} A \otimes \Lambda^m T^* M &\xrightarrow[\cong]{\varphi_\lambda} \Lambda^n \mathfrak{g} \otimes \Lambda^m TM \otimes \Lambda^m T^* M \xrightarrow[\cong]{\mathcal{K}} \Lambda^n \mathfrak{g} \\ \varepsilon \wedge \lambda X \otimes \varphi &\longleftarrow \varepsilon \otimes X \otimes \varphi \longmapsto \varepsilon \cdot \langle X, \varphi \rangle. \end{aligned}$$

Taking a local basis $\sigma_1, \dots, \sigma_n$ of \mathfrak{g} , W_1, \dots, W_m of TM and the duals W_1^*, \dots, W_m^* we see that $\bar{\mathcal{K}}(\sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^*) = \sigma_1 \wedge \dots \wedge \sigma_n$. To prove our lemma it is necessary to show the compatibility $D \bar{\mathcal{K}} \sim ad_A^n$ on these nonsingular sections only, i.e.

$$(ad_A^n)_\gamma(\sigma_1 \wedge \dots \wedge \sigma_n) = \bar{\mathcal{K}}(D_\gamma(\sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^*))$$

Let $[\gamma, \sigma_i] = \sum_j a_i^j \cdot \sigma_j$, $[\gamma, \lambda W_j] = \sum_k \tilde{a}_j^k \cdot \sigma_k + \sum_r b_j^r \cdot \lambda W_r$, then $[\#_A(\gamma), W_j] = \sum_r b_j^r \cdot \lambda W_r$. The right side of the above equation is equal to

$$\begin{aligned} &\bar{\mathcal{K}} \left(\sum a_i^i \cdot \sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^* + \right. \\ &\quad \left. + \sigma_1 \wedge \dots \wedge \sigma_n \wedge \sum b_j^j \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^* + \right. \\ &\quad \left. + \sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes \left(- \sum b_j^j \right) W_1^* \wedge \dots \wedge W_m^* \right) \\ &= \bar{\mathcal{K}} \left(\sum a_i^i \cdot \sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^* \right) \\ &= \sum a_i^i \cdot \left(\bar{\mathcal{K}}(\sigma_1 \wedge \dots \wedge \sigma_n \wedge \lambda W_1 \wedge \dots \wedge \lambda W_m \otimes W_1^* \wedge \dots \wedge W_m^*) \right) \\ &= (ad_A^n)_\gamma \cdot \sigma_1 \wedge \dots \wedge \sigma_n. \end{aligned}$$

Conclusion 7.12. (1)

$$H_{\partial_A, c}^{m+n}(A) \neq 0 \iff (\Lambda^n \mathfrak{g} \otimes or(M), ad_A^n \otimes \partial_A^{or}) \sim (M \times \mathbb{R}, \partial_A),$$

(2)

$$H_{\partial_A^{or},c}^{m+n}(A, or(M)) \neq 0 \iff (\Lambda^n \mathbf{g} \otimes or(M), ad_A^m \otimes \partial_A^{or}) \sim (or(M), \partial_A^{or}).$$

The following proposition generalizes Proposition 2.2. The proof is analogous.

Proposition 7.13. *For a representation $\nabla : A \rightarrow A(\xi)$ in a line vector bundle ξ the following conditions are equivalent:*

- (a) $(\xi, \nabla) \sim (or(M), \partial_A^{or})$,
- (b) $(\xi \otimes or(M), \nabla \otimes \partial_A^{or}) \sim (M \times \mathbb{R}, \partial_A)$.

In the sequel we need the notion of a modular class of a Lie algebroid [19], [4]. Firstly, we recall the characteristic classes of a representation $\nabla : A \rightarrow A(\xi)$ in a line vector bundle ξ . If ξ is trivial as a line bundle and $s \in \Gamma(\xi)$ is a nonsingular section of ξ we define a 1-cocycle $\theta \in \Omega^1(A)$ with respect to d_A defined by $\nabla_\gamma \nu = \theta_s(\gamma) \cdot s$. The class $\theta_\nabla = [\theta] \in H^1(A)$ is independent on the choice of s and is called *characteristic class* of A associated to the representation ∇ . For a general ξ , we define $\theta_\nabla = \frac{1}{2}\theta_{\nabla \otimes \nabla}$ ($\nabla \otimes \nabla$ is a flat representation in trivial line bundle $\xi \otimes \xi$). We add that if ξ is trivial, the last equation holds.

For next propositions and theorems we need the following lemma.

Lemma 7.14. *If ξ is a line bundle and $\{\varphi_\alpha\}$ is a collection of local trivialiations with the transition functions $\lambda_{\alpha\beta} : U_\alpha \times U_\beta \rightarrow \mathbb{R}$, $\varphi_\beta = \varphi_\alpha \cdot \lambda_{\alpha\beta}$, then there exist functions $f_\alpha > 0$ such that the local trivializations*

$$\bar{\varphi}_\alpha = \varphi_\alpha \cdot f_\alpha$$

($\bar{\varphi}_{\alpha,x} = \varphi_{\alpha,x} \cdot f_\alpha(x)$) have transition functions $\bar{\lambda}_{\alpha\beta} = \text{sgn } \lambda_{\alpha\beta}$.

In conclusion, each line bundle ξ possesses a system of local trivializations with transition functions equaling to ± 1 and then a family $\{s_\alpha\}$ of nonsingular \pm sections i.e. with transition functions equaling just to ± 1 .

Proof. Consider a line bundle ξ with a collection of local trivialiations $\{\varphi_\alpha\}$ and transition functions $\lambda_{\alpha\beta}$. The tensor product $\xi \otimes \xi$ is a trivialisable vector bundle with local system of trivializations $\{\varphi_\alpha \otimes \varphi_\alpha\}$. Choice a global trivialization $\rho : \xi \otimes \xi \rightarrow M \times \mathbb{R}$ such that $\rho_\alpha := \rho(\varphi_\alpha \otimes \varphi_\alpha(1 \otimes 1)) > 0$. We put

$$f_\alpha = \frac{1}{\sqrt{\rho_\alpha}} > 0.$$

We show that $\{f_\alpha\}$ is a required family. The transition functions $\bar{\lambda}_{\alpha\beta}$ for the collection of new local trivializations $\{\bar{\varphi}_\alpha := \varphi_\alpha \cdot f_\alpha\}$ are equal to $\bar{\lambda}_{\alpha\beta} = \lambda_{\alpha\beta} \cdot \frac{f_\beta}{f_\alpha}$ so that $\text{sgn } \bar{\lambda}_{\alpha\beta} = \text{sgn } \lambda_{\alpha\beta}$. On the other hand, $\rho_\beta = \rho(\varphi_\beta \otimes \varphi_\beta(1 \otimes 1)) = \lambda_{\alpha\beta}^2 \cdot \rho_\alpha$ so $\bar{\lambda}_{\alpha\beta}^2 = \left(\lambda_{\alpha\beta} \cdot \frac{f_\beta}{f_\alpha}\right)^2 = \lambda_{\alpha\beta}^2 \cdot \frac{\rho_\alpha}{\rho_\beta} = 1$ and next $|\bar{\lambda}_{\alpha\beta}| = 1$. Finally $\bar{\lambda}_{\alpha\beta} = \text{sgn } \bar{\lambda}_{\alpha\beta} = \text{sgn } \lambda_{\alpha\beta}$.

Lemma 7.15. *For an arbitrary line bundle ξ the characteristic class θ_∇ of a representation $\nabla : A \rightarrow A(\xi)$ can be computed via any family of local nonsingular \pm sections $\{s_\alpha\}$ (see Lemma 7.14) of ξ in the following way: the 1-differential A -form $\theta \in \Omega^1(A)$ defined by*

$$\theta(\gamma)|_{U_\alpha} s_\alpha = \nabla_\gamma(s_\alpha)$$

is a correctly defined d_A -cocycle and its cohomology class is equal to θ_∇ . $\theta_\nabla = 0$ if and only if there exists a family of local nonsingular ∇ -constant \pm sections $\{s_\alpha\}$, $\nabla s_\alpha = 0$. For transitive Lie algebroid A if $\theta_\nabla = 0$ then $\nabla^+ = 0$, so the isotropy Lie algebras \mathfrak{g}_x are unimodular.

The simple proof will be omitted.

We have $\theta_{\partial_A^{or}} = 0$. The modular class of a Lie algebroid A is by definition the characteristic class θ_A of the associated representation $D : A \rightarrow A(Q_A)$. According to Lemma 7.11 for a transitive Lie algebroid A we have $\theta_A = \theta_{ad_A^n}$ where $ad_A^n : A \rightarrow A(\Lambda^n \mathfrak{g})$ is the representation induced by the adjoint one $ad_A : A \rightarrow A(\mathfrak{g})$.

For real coefficients we have the following characterization of Lie algebroids in the case of the nontriviality of the top group of cohomology. Let $\{(U_\alpha, x_\alpha)\}$ be a coordinate open cover for the manifold M , with transition functions $g_{\alpha\beta} = x_\alpha \circ x_\beta^{-1}$. Each map x_α determines canonically a local trivialization \bar{x}_α of the line bundle $\Lambda^m TM$ and the family $\{\bar{x}_\alpha\}$ has the transition functions $J(g_{\alpha\beta})$.

Theorem 7.16. *The following conditions are equivalent*

- $H_{\partial_A, c}^{m+n}(A) \neq 0$,
- $H_{\partial_A, c}^{m+n}(A) \cong \mathbb{R}$ and $H(A)$ is a Poincaré algebra, i.e. the pairing $H^j(A) \times H_c^{m+n-j}(A) \rightarrow H_c^{m+n}(A) \cong \mathbb{R}$ is nondegenerate, $H^j(A) \cong (H_c^{m+n-j}(A))^*$,
- $(Q_A^{or}, D^{or}) \sim (M \times \mathbb{R}, \partial_A)$,
- $(\Lambda^n \mathfrak{g} \otimes or(M), ad_A^n \otimes \partial_A^{or}) \sim (M \times \mathbb{R}, \partial_A)$,
- $(\Lambda^n \mathfrak{g}, ad_A^n) \sim (or(M), \partial_A^{or})$, that is the holonomy homomorphism of $(\Lambda^n \mathfrak{g}, ad_A^n)$ is the same as for the orientation bundle $(or(M), \partial_A^{or})$.
- A is orientable vector bundle and $\theta_A = 0$ (in particular, \mathfrak{g}_x are unimodular).

Proof. (a) \iff (b) \iff (c) follows immediately from Theorem 7.10 for

$$(\xi, \nabla) = (M \times \mathbb{R}, \partial_A),$$

(c) \iff (d) by Lemma 7.11,

(d) \iff (e) see Proposition 7.13,

(e) \implies (f) indeed, $\theta_A = \theta_{ad_A^n} = \theta_{\partial_A^{or}} = 0$. The bundle $\Lambda^{m+n} A \cong \Lambda^n \mathfrak{g} \otimes \Lambda^m TM \cong or(M) \otimes \Lambda^m TM$ is trivial line bundle because it possesses a local system of trivializations with positive transition functions $|J(g_{\alpha\beta})|$.

(f) \implies (e) It is necessary to find a local system of nonsingular ad_A^n -constant sections $\{\sigma_\alpha\}$ of $\Lambda^n \mathfrak{g}$ with transition functions $\text{sgn } J(g_{\alpha\beta})$.

Fix a system of local trivializations $\{\psi_\alpha\}$ of $\Lambda^{m+n} A$ with positive transition functions $\gamma_{\alpha\beta} > 0$. We can choose a system of local trivializations $\{\varphi_\alpha\}$ of the line bundle $\Lambda^n \mathfrak{g}$

in such a way that $\varphi_\alpha \otimes \bar{x}_\alpha$ form a system of local trivializations of the line bundle $\Lambda^{m+n}A \cong \Lambda^n \mathfrak{g} \otimes \Lambda^m TM$ compatible with $\{\psi_\alpha\}$, i.e. such that

$$\varphi_\alpha \otimes \bar{x}_\alpha = g_\alpha \cdot \psi_\alpha, \quad g_\alpha > 0.$$

This implies that the transition functions $\lambda_{\alpha\beta}$ of the system $\{\varphi_\alpha\}$ have the sign of $J(g_{\alpha\beta})$. Indeed,

$$\varphi_\beta \otimes \bar{x}_\beta = \begin{cases} \varphi_\alpha \otimes \bar{x}_\alpha \cdot \lambda_{\alpha\beta} \cdot J(g_{\alpha\beta}) = \psi_\alpha \cdot g_\alpha \cdot \lambda_{\alpha\beta} \cdot J(g_{\alpha\beta}), \\ \psi_\beta \cdot g_\beta = \psi_\alpha \cdot \gamma_{\alpha\beta} \cdot g_\beta. \end{cases}$$

Therefore, since g_α, g_β and $\gamma_{\alpha\beta}$ are positive we have $\lambda_{\alpha\beta} \cdot J(g_{\alpha\beta}) > 0$, i.e.

$$\text{sgn } \lambda_{\alpha\beta} = \text{sgn } J(g_{\alpha\beta}).$$

By Lemma 7.14 there exists functions $f_\alpha > 0$ such that the local trivializations

$$\bar{\varphi}_\alpha = \varphi_\alpha \cdot f_\alpha$$

($\bar{\varphi}_{\alpha,x} = \varphi_{\alpha,x} \cdot f_\alpha(x)$) have transition functions $\bar{\lambda}_{\alpha\beta} = \text{sgn } \lambda_{\alpha\beta} = \text{sgn } J(g_{\alpha\beta})$. The family of \pm sections $\bar{\sigma}_\alpha = \bar{\varphi}_\alpha(1)$ determine a 1-cocycle $\theta \in \Omega^1(A)$ with respect to d_A defined by $\theta(\gamma)|_{U_\alpha} \cdot \bar{\sigma}_\alpha = (ad_A^n)_\gamma(\bar{\sigma}_\alpha)$ whose cohomology class is the characteristic class of the adjoint representation ad_A^n , $[\theta] = \theta_{ad_A^n}$. Since $\theta_{ad_A^n} = \theta_A = 0$ one has $\theta = d_A f$ for some function $f \in C^\infty(M)$, i.e.

$$\theta(\gamma) = (d_A f)(\gamma) = \partial_{\#_A(\gamma)} f.$$

Put $\sigma_\alpha = e^{-f} \cdot \bar{\sigma}_\alpha$. Then the transition functions of $\{\sigma_\alpha\}$ are equal to $\text{sgn } J(g_{\alpha\beta})$ and the sections σ_α are ad_A^n -constant.

In case of oriented manifold the above theorem yields:

Theorem 7.17. *If M is a oriented manifold then the following conditions are equivalent*

- (a) $H_{\partial_A, c}^{m+n}(A) \neq 0$,
- (b) $H_{\partial_A, c}^{m+n}(A) \cong \mathbb{R}$ and $H(A)$ is a Poincaré algebra, i.e. the pairing $H^j(A) \times H_c^{m+n-j}(A) \rightarrow H_c^{m+n}(A) \cong \mathbb{R}$ is nondegenerate, $H^j(A) \cong (H_c^{m+n-j}(A))^*$,
- (c) $(\Lambda^n \mathfrak{g}, ad_A^n) \sim (M \times \mathbb{R}, \partial_A)$, i.e. there exists a global nonsingular section $\varepsilon \in \Gamma(\Lambda^n \mathfrak{g})$ which is ad_A^n -constant, that is, A is a TUIO-Lie algebroid, see [13],
- (d) \mathfrak{g} is orientable and $\theta_A = 0$.

The independent proof of the implication (c) \implies (b) one can be found in [13].

Finally we give a characterization of Lie algebroids whose the top group of cohomology with coefficients in the orientation bundle $or(M)$ is not trivial.

Theorem 7.18. *The following conditions are equivalent:*

- (a) $H_{\partial_A^{or}, c}^{m+n}(A, or(M)) \neq 0$,

(b) $H_{\partial_A^{or},c}^{m+n}(A, or(M)) \cong \mathbb{R}$ and the pairing

$$H^j(A) \times H_{\partial_A^{or},c}^{m+n-j}(A, or(M)) \rightarrow H_{\partial_A^{or},c}^{m+n}(A, or(M)) \cong \mathbb{R}$$

is not degenerate, i.e. $H^j(A) \cong (H_{\partial_A^{or},c}^{m+n-j}(A, or(M)))^*$,

(c) $(Q_A^{or}, D^{or}) \sim (or(M), \partial_A^{or})$,

(d) $(\Lambda^n \mathfrak{g} \otimes or(M), ad_A^n \otimes \partial_A^{or}) \sim (or(M), \partial_A^{or})$,

(e) $(\Lambda^n \mathfrak{g}, ad_A^n) \sim (M \times \mathbb{R}, \partial_A)$,

(f) \mathfrak{g} is orientable and there exists a global nonsingular section $\varepsilon \in \Gamma(\xi)$ which is ad_A^n -constant (i.e. A is a TUIO-Lie algebroid, see [13]),

(g) \mathfrak{g} is orientable and $\theta_A = 0$.

Proof. Only the implication (d) \implies (e) needs a proof. Since

$$(or(M) \otimes or(M), \partial_A^{or} \otimes \partial_A^{or}) \sim (M \times \mathbb{R}, \partial_A)$$

one has

$$\begin{aligned} (\Lambda^n \mathfrak{g}, ad_A^n) &\sim (\Lambda^n \mathfrak{g} \otimes or(M) \otimes or(M), ad_A^n \otimes \partial_A^{or} \otimes \partial_A^{or}) \\ &\stackrel{(d)}{\sim} (or(M) \otimes or(M), \partial_A^{or} \otimes \partial_A^{or}) \\ &\sim (M \times \mathbb{R}, \partial_A). \end{aligned}$$

For an orientable manifold we get Theorem 7.17.

7.5 Remarks on Example 5.3 from [4].

In the cited paper there is an example of nonregular Lie algebroid for which the E-L-W cohomological pairing is not necessary nondegenerate. In the text of Example 5.3 from [4] there are some inaccuracies (concerning dimensional of the group of cohomology) which we remove here. We prove additionally that there is no line representation for which the cohomological pairing is nondegenerate and we prove that the E-L-W representation is not exceptional. The example is the Lie transformation algebroid $A = \mathfrak{g} \times M \rightarrow M$ associated with the infinitesimal action $\gamma : \mathfrak{g} \rightarrow \mathfrak{X}(M)$ of a finitely dimensional Lie algebra \mathfrak{g} on a manifold M . The anchor is given by $\rho(v, x) = \gamma(v)_x$, and Lie bracket by

$$[[a, b]](x) = [a(x), b(x)] + \gamma(a(x))_x(b) - \gamma(b(x))_x(a),$$

$a, b \in C^\infty(M, \mathfrak{g}) \cong \Gamma(\mathfrak{g} \times M)$ and $x \in M$. The vector field $X = x^N \frac{d}{dx}$ on \mathbb{R} ($N \in \mathbb{N}$) defines an action of the 1-dimensional Lie algebra $\mathfrak{g} = \mathbb{R}$ on $M = \mathbb{R}$ by $\gamma : \mathbb{R} \rightarrow \mathfrak{X}(\mathbb{R})$, $\gamma(t) = t \cdot X$. Let A be the transformation Lie algebroid associated with γ . Then $\Gamma(A) = C^\infty(\mathbb{R})$, $\#_x : A_x = \mathbb{R} \rightarrow T_x M$, $t \mapsto t \cdot x^N \cdot \frac{d}{dx}$, $[[a, b]] = x^N \cdot (a \cdot b' - b \cdot a')$, $\Omega^0(A) = C^\infty(\mathbb{R})$,

$$\Omega^1(A) = \Gamma(A^*) = C^\infty(\mathbb{R}, \mathbb{R}^*) = \Omega^1(\mathbb{R}) \cong C^\infty(\mathbb{R}), \quad f dx \mapsto f, \quad (35)$$

and, clearly, $\Omega^2(A) = 0$.

Lemma 7.19. $H^1(A) \cong \mathbb{R}^N$.

Proof. By definition $d_A : C^\infty(\mathbb{R}) \rightarrow \Omega^1(A) \cong C^\infty(\mathbb{R})$, $d_A(f)(a) = \#(a)(f) = a \cdot x^N \cdot f'$, and therefore $d_A(f) = x^N \cdot f'$ and

$$H^1(A) = C^\infty(\mathbb{R}) / \{x^N \cdot f', f \in C^\infty(\mathbb{R})\} = C^\infty(\mathbb{R}) /_{x^N \cdot C^\infty(\mathbb{R})} \cong \mathbb{R}^N.$$

Indeed, the classes of functions x^0, x^1, \dots, x^{N-1} form a basis of $C^\infty(\mathbb{R}) /_{x^N \cdot C^\infty(\mathbb{R})}$ because the classes are linearly independent and for any $f \in C^\infty(\mathbb{R})$ the equality $[f] = \sum_{i=0}^{N-1} a_i [x^i]$ holds where $a_i = \frac{f^{(i)}(0)}{i!}$.

Proposition 7.20. For each linear representation $\nabla : A \rightarrow A(\xi)$ we have (1) $H^0_{\nabla,c}(A, \xi) = 0$, (2) $H^1_{\nabla,c}(A, \xi) \neq 0$.

Therefore, for each representation ∇ of A in a line bundle ξ the cohomological pairing $H^1(A) \times H^0_{\nabla,c}(A, \xi) \rightarrow H^1_{\nabla,c}(A, \xi)$ is not nondegenerate even in a weak manner[‡].

Proof. (1): The line bundle ξ over \mathbb{R} is trivial $\xi = M \times \mathbb{R}$ ($M = \mathbb{R}$) so each representation $\nabla : A \rightarrow A(\xi)$ is of the form $\nabla = \partial_A^\omega$ for some 1-form $\omega \in \Omega^1(A)$. Let $\omega(a) = g \cdot a$ for $g \in C^\infty(\mathbb{R})$. Then $(\partial_A^\omega)_a(f) = (\partial_A)_a(f) + \omega(a) \cdot f = a \cdot x^N \cdot f' + a \cdot g \cdot f$ and

$$H^0_{\nabla,c}(A, \xi) = \{f \in C_c^\infty(\mathbb{R}) ; x^N \cdot f' + g \cdot f = 0\} = 0.$$

by the uniqueness of the Cauchy problem for the differential equation $y' = -\frac{g(x)}{x^N} \cdot y$.

(2) $H^1_{\nabla,c}(A, \xi) = C_c^\infty(\mathbb{R}) /_{\{x^N \cdot f' + g \cdot f; f \in C_c^\infty(\mathbb{R})\}} \neq 0$. To prove this we find a compactly supported function $h \in C_c^\infty(\mathbb{R})$ such that the differential equation

$$x^N \cdot y' + g \cdot y = h \tag{36}$$

has no global solution $y \in C_c^\infty(\mathbb{R})$.

Case $g(0) = 0$. For any h such that $h(0) \neq 0$ there is no solution of (36).

Case $g(0) \neq 0$. Let $|g(x)| \geq \delta > 0$ for $|x| \leq \varepsilon$, $\varepsilon > 0$. Take any function $h \in C_c^\infty(\mathbb{R})$ such that $h \geq 0$, $h \neq 0$ and $\text{supp } h \subset [\alpha, \beta] \subset (\varepsilon, \infty)$. The elementary theory of linear differential equations [the formula solving the Cauchy problem in the form of denoted integrals] yields easily that no global compactly supported solution of (36) exists.

Consider the E-L-W representation $D : A \rightarrow A(Q_A)$. We see that $Q_A = A \otimes T^*\mathbb{R} \cong M \times \mathbb{R}$ [$M = \mathbb{R}$] so $\Gamma(Q_A) = \Gamma(A) \otimes \Omega^1(\mathbb{R}) \cong C^\infty(M)$ by $1 \otimes f dx \mapsto f$ and that D is equivalent to ∂_A^ω for $\omega \cong (x^N)'$ (with respect to isomorphism (35)). According to Proposition (7.20) the top group of cohomology of A for trivial and for E-L-W representations are nontrivial. We prove that this representations are not isomorphic so the E-L-W representation is not exceptional.

[‡] A pairing $F : V \times W \rightarrow U$ is called weakly non-degenerated if both null spaces $N_1 = \{v \in V; F(v, \cdot)\}$ and $N_2 = \{w \in W; F(\cdot, w)\}$ are zero

Proposition 7.21. *The A -flat line bundles $(M \times \mathbb{R}, \partial_A)$ and (Q_A, D) [$M = \mathbb{R}$] are not isomorphic.*

Proof. Let $\varphi : M \times \mathbb{R} \rightarrow Q_A$ be a linear homomorphism compatible with ∂_A and D . φ is of the form $\varphi(f) = 1 \otimes g \cdot f \cdot dx$ for some $g \in C^\infty(M)$. The equality $D_a(\varphi(f)) = \varphi((\partial_A)_a f)$ yields $a \cdot (x^N \cdot fg)' = a \cdot x^N \cdot f' \cdot g$, therefore $(x^N \cdot g)' = 0$ which produces $g \equiv 0$ and that φ is not an isomorphism.

References

- [1] R. Bott and L. Tu: *Differential forms in algebraic topology*, GTM 82, Springer-Verlag, 1982.
- [2] S.S. Chern, F. Hirzebruch and J-P. Serre: “On the index of a fibered manifold”, *Proc. AMS*, Vol. 8, (1957), pp. 587–596.
- [3] M. Crainic: “Differentiable and algebroid cohomology, Van Est isomorphisms, and characteristic classes”, preprint, arXiv:math.DG/0008064, *Commentarii Mathematici Helvetici*, to appear.
- [4] S. Evens, J-H. Lu and A. Weinstein: “Transverse measures, the modular class and a cohomology pairing for Lie algebroids”, *Quart. J. Math. Oxford*, Vol. 50, (1999), pp. 417–436.
- [5] R.L. Fernandes: “Lie algebroids, holonomy and characteristic classes”, preprint DG/007132, *Advances in Mathematics*, Vol. 170, (2002), pp. 119–179.
- [6] F. Guedira and A. Lichnerowicz: “Géométrie des algèbres de Lie locales de Kirillov”, *J. Math. Pures Appl.*, Vol. 63, (1984), pp. 407–484.
- [7] W. Greub, S. Halperin and R. Vanstone: *Connections, Curvature and Cohomology*, New York and London, Vol. I, 1971; Vol. II, 1973.
- [8] S. Haller and T. Rybicki: “Reduction for locally conformal symplectic manifolds”, *J. Geom. Phys.*, Vol. 37, (2001), pp. 262–271.
- [9] G. Hochschild and J.-P. Serre: “Cohomology of Lie algebras”, *Ann. Math.*, Vol. 57, (1953), pp. 591–603.
- [10] J. Huebschmann: “Duality for Lie-Rinehart algebras and the modular class”, *J. Reine Angew. Math*, Vol. 510, (1999), pp. 103–159.
- [11] V. Itskov, M. Karashev and Y. Vorobjev: “Infinitesimal Poisson Cohomology”, *Amer. Math. Soc. Transl. (2)*, Vol. 187, (1998).
- [12] J. Kubarski: “Invariant cohomology of regular Lie algebroids”, In: X. Masa, E. Macias-Virgos, J. Alvarez Lopez (Eds.): *Proceedings of the VII International Colloquium on Differential Geometry ANALYSIS AND GEOMETRY IN FOLIATED MANIFOLDS*, Santiago de Compostela, Spain, July 1994, World Scientific, Singapore, 1995, pp. 137–151.
- [13] J. Kubarski: “Fibre integral in regular Lie algebroids”, In: *Proceedings of the Conference: New Developments in Differential Geometry, Budapest 1996*, Budapest, Hungary, 27-30 July 1996, Kluwer Academic Publishers, 1999.
- [14] J. Kubarski: “Poincaré duality for transitive unimodular invariantly oriented Lie algebroids”, *Topology and Its Applications*, Vol. 121(3), (2002), pp. 333–355.

-
- [15] R. Kadobianski, J. Kubarski, V. Kushnirevitch and R. Wolak: “Transitive Lie algebroids of rank 1 and locally conformal symplectic structures”, *Journal of Geometry and Physics*, Vol. 46, (2003), pp. 151–158.
- [16] J. Kubarski and A.S. Mishchenko: “Lie algebroids: spectral sequences and signature”, *Sbornik: Mathematics*, Vol. 194(7), (2003), pp. 1079–1103.
- [17] K. Mackenzie: *Lie Groupoids and Lie Algebroids in Differential Geometry*, London Math. Soc., Lecture Notes, Series 124, Cambridge University Press, 1987.
- [18] M. Spivak: *A Comprehensive Introduction to Differential Geometry*, Vol. I, 2nd Ed., Publish or Perish Inc., Berkeley, 1979.
- [19] A. Weinstein: “The modular automorphism group of a Poisson manifold”, *J. Geom. Phys.*, Vol. 23, (1997), pp. 379–394.