

Ideals with bases of unbounded Borel complexity

Piotr Borodulin–Nadzieja*, Szymon Głąb

Abstract

We present several naturally defined σ -ideals which have Borel bases but, unlike for the classical examples, these bases are not of bounded Borel complexity. We investigate set-theoretic properties of such σ -ideals.

1 Introduction

Consider the σ -ideal of subsets of the plane consisting of sets which for all $\varepsilon > 0$ can be covered by an open set whose all vertical sections have measure less than ε . It seems natural to suppose that this ideal has similar properties to those of the ideal of null sets, e.g. Gabriel Mokobodzki conjectured that under Martin’s Axiom the additivity of this ideal equals continuum (see [9], Problem 32Rd). It turned out to be not true. Cichoń and Pawlikowski proved in [6] that the additivity of this ideal, called there Mokobodzki ideal, equals ω_1 in ZFC.

In the same paper Cichoń and Pawlikowski considered also σ -ideals of subsets of $[0, 1]^2$ which can be covered by Borel sets whose every vertical section is small (i.e. of Lebesgue measure 0 or meager). These ideals also have additivity ω_1 . Cichoń and Pawlikowski observed one more interesting property: these ideals do not have Borel bases of bounded Borel complexity (although, clearly, they have Borel bases). In other words, $\mathcal{M}_\alpha \subsetneq \mathcal{M}$ for each α , where \mathcal{M} is such an ideal and \mathcal{M}_α is the σ -ideal generated by $\mathcal{M} \cap \Sigma_\alpha^0$.

In this paper we consider certain modifications of Mokobodzki ideals, e.g. the σ -ideal of subsets of the plane which are small not only on vertical sections but also on horizontal ones or which are small *in every direction*. In Section 3 we prove that such σ -ideals also do not have Borel bases of bounded Borel complexity.

A set A is in Mokobodzki ideal if its *every* section is small. One can ask if it is possible to change *every* to *almost every* and still having a σ -ideal with bases of unbounded Borel complexity. It can be done if we interpret the word “almost” correctly. Notice that ideals of sets whose almost every (with respect to Lebesgue measure or with respect to Baire category) section is small (Lebesgue null or meager, respectively) are the ideals of null or meager subsets of the real plane, and therefore they have bases of bounded Borel complexity (e.g. consisting of Π_2^0 or Σ_2^0 sets, respectively). However, in Section 4

*The first author was partially supported by the Polish Ministry of Science and Higher Education under grant no. N N201 418939.

2010 *Mathematics Subject Classification*: Primary 54A35, 28A05.

Keywords: ideals on the plane, Mokobodzki ideal, Fubini products, uniformization, property (M).

we proved that *every* can be changed to *almost every with respect to a σ -ideal having property (M)*.

In Section 5 we investigate set-theoretic properties of the σ -ideals considered in the paper. In the last section we present a remark concerning ideals with property (M) and we state some open questions. Facts proved in the two last sections indicate that Mokobodzki ideals and other ideals considered here differ from *Null* and *Meager* in many aspects.

Acknowledgements. The authors would like to thank Jacek Cichoń for introducing them to the notion of Mokobodzki ideal, Piotr Zakrzewski for valuable suggestions, and an anonymous referee for the careful examination of the paper and for helpful comments.

2 Preliminaries

All terminology which is not explained here, can be found e.g. in [11] and [5]. By $\Sigma_\alpha^0(X)$, $\Pi_\alpha^0(X)$, $\text{Borel}(X)$, $\Sigma_1^1(X)$ and $\Pi_1^1(X)$ we mean the families of, respectively, Σ_α^0 , Π_α^0 , Borel, analytic and coanalytic subsets of a Polish space X . Usually, X will be known from the context and we omit it. By $\pi_1: X \times Y \rightarrow X$ and $\pi_2: X \times Y \rightarrow Y$ we denote the projections on the first and on the second coordinates, i.e. $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$ for every $(x, y) \in X \times Y$. For $A \subseteq X \times Y$ define a vertical section of A at a point x as $A_x = \{y \in Y: (x, y) \in A\}$ and a horizontal section of A at a point y as $A^y = \{x \in X: (x, y) \in A\}$. In the paper we use several times the fact that if A and f are Borel and $f|A$ is injective, then $f(A)$ is Borel (see [11, 15.1]).

Let \mathcal{I} be a σ -ideal on Y and \mathcal{J} be a σ -ideal on X . The Fubini product $\mathcal{J} \otimes \mathcal{I}$ is the σ -ideal on $X \times Y$ generated by the family $\{B \in \text{Borel}(X \times Y): \{x: B_x \notin \mathcal{I}\} \in \mathcal{J}\}$. We say that a σ -ideal \mathcal{I} is Σ_α^0 -on- Π_α^0 if $\{x: B_x \in \mathcal{I}\} \in \Pi_\alpha^0$ for every $B \in \Sigma_\alpha^0(X \times X)$. Let A be a Σ_α^0 subset of $[0, 1]^2$. By [11, 22.22] the set $\{x: A_x \text{ is non-meager}\}$ is Σ_α^0 . Then $\{x: A_x \text{ is meager}\} = \{x: A_x \text{ is non-meager}\}^c$ is Π_α^0 . Using [11, 22.25] we obtain that $\{x: A_x \text{ is not null}\}$ is Σ_α^0 . Hence $\{x: A_x \text{ is null}\}$ is Π_α^0 . This shows that the ideals *Meager* and *Null* are Σ_α^0 -on- Π_α^0 for every $\alpha < \omega_1$. In a similar way we define properties Borel-on-Borel, Π_1^1 -on- Σ_1^1 etc. Note that the ideals *Meager* and *Null* are Π_1^1 -on- Σ_1^1 (see [11, 29.22 and 29.26]).

Let \mathcal{J} be a σ -ideal. A family $\mathcal{B} \subseteq \mathcal{J}$ is called a base of \mathcal{J} if any set of \mathcal{J} is contained in some set from \mathcal{B} . If there is a base of \mathcal{J} consisting of Borel or Σ_α^0 sets, then we say that \mathcal{J} has a Borel or Σ_α^0 base, respectively. For an ordinal number $\alpha \leq \omega_1$, by \mathcal{J}_α we denote the σ -ideal generated by $\mathcal{J} \cap \Sigma_\alpha^0$. Note that if \mathcal{J} has a Borel base, then $\mathcal{J} = \bigcup_{\alpha < \omega_1} \mathcal{J}_\alpha$. Let \mathcal{I} be a σ -ideal of subsets of an uncountable Polish space X . We say that $A \subseteq X^2$ is in the σ -ideal $\mathcal{M}(\mathcal{I})$ if there is a Borel set $B \supseteq A$ such that $B_x \in \mathcal{I}$ for every $x \in X$. We write $\mathcal{M}_\alpha(\mathcal{I})$ instead of $(\mathcal{M}(\mathcal{I}))_\alpha$. We will say that a σ -ideal \mathcal{J} with a Borel base has *the complex Borel base property* if for every $\alpha < \omega_1$ we have $\mathcal{J}_\alpha \subsetneq \mathcal{J}$. In [6] the authors mentioned that $\mathcal{M}(\text{Null})$ and $\mathcal{M}(\text{Meager})$ have the complex Borel base property.

For a σ -ideal \mathcal{I} of subsets of X^2 and a family of functions $\mathcal{F} \subseteq X^X$ define a σ -ideal $\mathcal{M}(\mathcal{F}, \mathcal{I})$ on X^2 in the following way: $Y \subseteq X^2$ belongs to $\mathcal{M}(\mathcal{F}, \mathcal{I})$ whenever $Y \in \mathcal{M}(\mathcal{I})$ and Y can be covered by a Borel set $B \subseteq X^2$ such that $\{x: (x, f(x)) \in B\} \in \mathcal{I}$ for every $f \in \mathcal{F}$. As before, $\mathcal{M}_\alpha(\mathcal{F}, \mathcal{I}) = (\mathcal{M}(\mathcal{F}, \mathcal{I}))_\alpha$. Note that if $\mathcal{F} = \emptyset$, then $\mathcal{M}(\mathcal{F}, \mathcal{I}) = \mathcal{M}(\mathcal{I})$; if \mathcal{F} consists of all constant functions $f \equiv y$, then $\{x: (x, f(x)) \in$

$B\} = \{x: (x, y) \in B\} = B^y$ which means that all horizontal sections B^y of B are in \mathcal{I} ; if \mathcal{F} is a family of all linear mappings, then $\mathcal{M}(\mathcal{F}, \mathcal{I})$ consists of all subsets of real plane which can be covered by Borel set \mathcal{I} -small in every direction, etc. Hence, $\mathcal{M}_\alpha(\mathcal{F}, \mathcal{I})$ seems to be a natural generalization of $\mathcal{M}_\alpha(\mathcal{I})$. In Section 3 we discuss the properties of such σ -ideals and we show that, under natural assumptions on \mathcal{F} , they have the complex Borel base property.

One of the well-known uniformization theorems states that if every nonempty section of a Borel set $A \subseteq X^2$ is not in \mathcal{I} (where X is an uncountable Polish space and \mathcal{I} is a Borel-on-Borel ideal), then A has a Borel uniformization, i.e. a Borel function $f: \pi_1[A] \rightarrow X$ such that $f(x) \in A_x$ for every $x \in \pi_1[A]$ [11, 18.6]. Recently, Petr Holický proved a theorem which gives an information about the Borel class of this uniformization. Here we state a simplified version of Holický's result needed for our purposes. It immediately follows from [10, Theorem 3.3] and [10, Theorem 3.4]. Recall that a function $f: X \rightarrow Y$ is Σ_α^0 -measurable if for every $E \in \Sigma_1^0(Y)$, the set $f^{-1}[E] \in \Sigma_\alpha^0(X)$. The graph of Σ_α^0 -measurable function belongs to $\Pi_\alpha^0(X \times Y)$ if X and Y are Polish (see, e.g. [12] §31, VII, Thm 1).

Theorem 2.1 (Holický) *Suppose X is an uncountable Polish space. Let \mathcal{I} be a σ -ideal of subsets of X which is Σ_α^0 -on- Π_α^0 for some $2 \leq \alpha < \omega_1$ and which contains all singletons. Let $A \subseteq X^2$ be such that $A_x \notin \mathcal{I}$ for every $x \in \pi_1[A]$. If A is of class Σ_α^0 , then there is a Borel function $F: \{0, 1\}^\omega \times X \rightarrow X$ such that:*

$$\begin{aligned} \forall x \in \{0, 1\}^\omega \quad (y \mapsto F(x, y) \text{ is a } \Sigma_\alpha^0\text{-measurable uniformization of } A) \\ \forall y \in X \quad (x \mapsto F(x, y) \text{ is continuous and 1-1). \end{aligned}$$

In particular, there is a Σ_α^0 -measurable uniformization of A .

The following theorem was proved in [6]. We repeat here its proof since we slightly modify its conclusion and we will use a similar argument later.

Theorem 2.2 (Cichoń, Pawlikowski, Lemma 2.3 in [6]) *Assume \mathcal{I} is a σ -ideal of subsets of an uncountable Polish space X such that $X \notin \mathcal{I}$. For every $\alpha < \omega_1$ there is a Π_α^0 set $A \subseteq X^2$ such that for every $M \in \mathcal{M}_\alpha(\mathcal{I})$ there is $x \in X$ such that $\emptyset \neq A_x \subseteq M_x^c$. If, additionally, \mathcal{I} is Σ_α^0 -on- Π_α^0 , then we can assume that $A_x^c \in \mathcal{I}$ for every $x \in \pi_1[A]$.*

Proof. Let $\alpha < \omega_1$. By [11, 22.3] there is a universal set $U \subseteq X \times X^2$ for the pointclass $\Sigma_\alpha^0(X^2)$, i.e. for every $E \subseteq X^2$ such that $E \in \Sigma_\alpha^0$ there is $x \in X$ with $E = U_x$. Put $A = \{(x, y) \in X^2: (x, x, y) \notin U\}$. Clearly, A is a Π_α^0 subset of X^2 . Let $M \in \mathcal{M}_\alpha(\mathcal{I})$. There is $E \in \Sigma_\alpha^0 \cap \mathcal{M}_\alpha(\mathcal{I})$ such that $M \subseteq E$. Since U is universal for $\Sigma_\alpha^0(X^2)$, there is $x_0 \in X$ with $E = U_{x_0}$. Since $A_{x_0} = X \setminus U_{(x_0, x_0)} = X \setminus (U_{x_0})_{x_0} = E_{x_0}^c$, then $A_{x_0}^c \in \mathcal{I}$.

Now assume that \mathcal{I} is Σ_α^0 -on- Π_α^0 . Then the set $\{x: A_x^c \in \mathcal{I}\}$ is Π_α^0 . Define

$$A' = A \cap (\{x: A_x^c \in \mathcal{I}\} \times X).$$

Of course A' is Π_α^0 . Fix $x \in X$ such that $U_x \in \mathcal{M}_\alpha(\mathcal{I})$ and notice that $A_x = \{y: (x, x, y) \notin U\}$ and $U_{(x, x)} \in \mathcal{I}$. Therefore $A_x^c \in \mathcal{I}$ and $A_x = A'_x$. Consequently, for any $M \in \mathcal{M}_\alpha(\mathcal{I})$ there is $x \in X$ such that $A'_x \subseteq M_x^c$. Thus, we can assume without loss of generality that $A_x^c \in \mathcal{I}$ for every $x \in \pi_1[A]$. ■

3 Ideals $\mathcal{M}(\mathcal{F}, \mathcal{I})$

In this section (X, \cdot) stands for an uncountable Polish group. One can e.g. think about the group $X = [0, 1)$ with the addition modulo 1 or $X = 2^\omega$ with the standard additive operation. Let \mathcal{I} be either the σ -ideal of meager subsets of X or a σ -ideal of null subsets of X with respect to a right-invariant σ -finite measure on X .

We do not consider \mathcal{I} in more general setting, since we will need many particular properties of the above ideals: Fubini property, ccc, right-invariance, Σ_α^0 -on- Π_α^0 for each $\alpha < \omega_1$. Zakrzewski's result from [15] implies that in the case $X = 2^\omega$ the ideals $\mathcal{N}ull$ and $\mathcal{M}eager$ are the only σ -ideals satisfying the above properties. This and other well-known results (see [8]) suggest that in case of arbitrary Polish group, if \mathcal{I} satisfy the above properties then it is at least isomorphic to one of the above ideals.

If \mathcal{I} is an ideal of null sets, then we will additionally assume that the measure is σ -finite, since we will need the following property: every Borel function $f: B \rightarrow X$, where $B \subseteq X$ is \mathcal{I} -positive and Borel, is continuous on an \mathcal{I} -positive Σ_3^0 subset of B . This property is satisfied under the above condition (see [11], Thm. 17.12) and in the case \mathcal{I} is the ideal of meager subsets.

For the functions $f, g: X \rightarrow X$ let $(f \cdot g)(x) = f(x) \cdot g(x)$.

A family of functions $\mathcal{F} \subseteq X^X$ is *ubiquitous with respect to an ideal \mathcal{I}* (or \mathcal{I} -ubiquitous) if for every Borel function $g: X \rightarrow X$ there is a Borel set $B \notin \mathcal{I}$ and a function $f \in \mathcal{F}$ such that $f|_B = g|_B$. The family of continuous functions is a natural example of $\mathcal{N}ull$ - and $\mathcal{M}eager$ -ubiquitous family (it follows from Luzin Theorem [11, 17.12] and Nikodym Theorem [11, 8.38]). On the other hand, there are families of Borel functions $f: [0, 1) \rightarrow [0, 1)$ which are closed under the addition modulo 1 but are not ubiquitous neither with respect to $\mathcal{N}ull$ nor to $\mathcal{M}eager$ ideals: the empty family, the constant functions, the linear functions. Note that also the family of polynomials is not ubiquitous neither with respect to $\mathcal{N}ull$ nor to $\mathcal{M}eager$: e.g. the exponential function cannot equal to a polynomial on a set with an accumulation point (as zeros of a holomorphic function must be isolated).

We will show that if a family \mathcal{F} of Borel functions is left shift invariant and is not \mathcal{I} -ubiquitous, then any Π_α^0 set with large sections and a big projection on x -axis have a uniformization with a graph of class $\Sigma_{\alpha+2}^0$ witnessing that \mathcal{F} is not \mathcal{I} -ubiquitous.

Lemma 3.1 *Assume that A is a Borel subset of X^2 such that $A_x \notin \mathcal{I}$ for every $x \in \pi_1[A] \notin \mathcal{I}$. For each Borel mapping $h: X \rightarrow X$ we can find a Borel set $B \subseteq \pi_1[A]$ with $B \notin \mathcal{I}$ and $y \in X$ such that $y \cdot h(x) \in A_x$ for every $x \in B$.*

Proof. Define $\varphi: X^2 \rightarrow X^2$ by $\varphi(x, y) = (x, y \cdot (h(x))^{-1})$. Then for every x

$$(\varphi[A])_x = (\{(x, y \cdot (h(x))^{-1}): (x, y) \in A\})_x = \{y \cdot (h(x))^{-1}: y \in A_x\} = A_x \cdot (h(x))^{-1}.$$

Since \mathcal{I} is right invariant, then

$$(\varphi[A])_x \in \mathcal{I} \text{ iff } A_x \in \mathcal{I}$$

and, thus, $(\varphi[A])_x \notin \mathcal{I}$ for every $x \in \pi_1[A] \notin \mathcal{I}$. The set $\varphi[A]$ is Borel because φ is a Borel one-to-one mapping. By the Fubini property of \mathcal{I} there is $y \in X$ such that

$$B = (\varphi[A])^y \in \text{Borel} \setminus \mathcal{I}.$$

Let $x \in B$. Then $(x, y) \in \varphi[A]$ and, therefore, $(x, y \cdot h(x) \cdot (h(x))^{-1}) \in \varphi[A]$ but this means that $(x, y \cdot h(x)) \in A$. So, $y \cdot h(x) \in A_x$ for every $x \in B$. ■

Lemma 3.2 *Let $3 \leq \alpha < \omega_1$. Assume $A \subseteq X^2$ is a Π_α^0 set such that $A_x \notin \mathcal{I}$ for every $x \in \pi_1[A] \notin \mathcal{I}$. Then, for a Borel function $h: X \rightarrow X$ there is a countable collection \mathcal{B} of pairwise disjoint Borel sets and a uniformization $f: \pi_1[A] \rightarrow X$ of A such that*

(i) $\pi_1[A] \setminus \bigcup \mathcal{B} \in \mathcal{I}$,

(ii) for every $B \in \mathcal{B}$ there is $y_B \in X$ such that $f|_B = y_B \cdot h$,

(iii) f is $\Sigma_{\alpha+1}^0$ -measurable (and, consequently, its graph is $\Pi_{\alpha+1}^0$).

Proof. Fix $A \subseteq X^2$ and $h: X \rightarrow X$ as above. Since $A \in \Sigma_{\alpha+1}^0$, we can use Theorem 2.1 to fix a $\Sigma_{\alpha+1}^0$ -measurable uniformization $g: \pi_1[A] \rightarrow X$ of A . Use Lemma 3.1 to find a Borel set B_0 and y_0 such that $B_0 \subseteq \pi_1[A]$, $B_0 \notin \mathcal{I}$ and $y_0 \cdot h(x) \in A_x$ for each $x \in B_0$. We can assume that $(y_0 \cdot h)|_{B_0}$ is continuous and B_0 is Σ_3^0 (more precisely, Π_2^0 if $\mathcal{I} = \text{Meager}$, and Σ_2^0 if $\mathcal{I} = \text{Null}$), shrinking B_0 if needed.

Assume now that we have constructed a family of pairwise disjoint Σ_3^0 sets $\{B_\xi: \xi < \beta\}$ and a family of points $\{y_\xi: \xi < \beta\}$ such that $B_\xi \subseteq \pi_1[A]$, $B_\xi \notin \mathcal{I}$, $y_\xi \cdot h(x) \in A_x$ for each $x \in B_\xi$, $(y_\xi \cdot h)|_{B_\xi}$ is continuous and B_ξ is Σ_3^0 .

If $Y = \pi_1[A] \setminus \bigcup_{\xi < \beta} B_\xi \notin \mathcal{I}$, then use Lemma 3.1 to find a Borel set B_β and a point y_β such that $B_\beta \subseteq \pi_1[A \cap (Y \times X)] = Y \cap \pi_1[A]$, $B_\beta \notin \mathcal{I}$, $y_\beta \cdot h(x) \in A_x$ for each $x \in B_\beta$, $(y_\beta \cdot h)|_{B_\beta}$ is continuous and B_β is Σ_3^0 . Since \mathcal{I} is ccc, there is $\beta < \omega_1$ such that $\pi_1[A] \setminus \bigcup_{\xi < \beta} B_\xi \in \mathcal{I}$. Let $\mathcal{B} = \{B_\xi: \xi < \beta\}$. Define $f: \pi_1[A] \rightarrow X$ in the following way. Let $f(x) = y_\xi \cdot h(x)$ if $x \in B_\xi$ and let $f(x) = g(x)$ for $x \in \pi_1[A] \setminus \bigcup \mathcal{B}$.

We have to verify that f defined in this way is $\Sigma_{\alpha+1}^0$ -measurable. Indeed, for every $\xi < \beta$ the function $h_\xi: B_\xi \rightarrow X$ defined by $h_\xi(x) = y_\xi \cdot h(x)$ is continuous on B_ξ . So, for $E \in \Sigma_1^0$ we have

$$f^{-1}[E] = \bigcup_{\xi < \beta} (h_\xi^{-1}[E] \cap B_\xi) \cup g^{-1}[E]$$

which is a countable union of $\Sigma_{\alpha+1}^0$ sets, so a $\Sigma_{\alpha+1}^0$ set. ■

Now we will use the above lemmas to prove that $\mathcal{M}(\mathcal{F}, \mathcal{I})$ has the complex base property under certain assumptions on \mathcal{F} .

Theorem 3.3 *Let $\mathcal{F} \subseteq X^X$ be a family of Borel functions which is not \mathcal{I} -ubiquitous. Assume that \mathcal{F} is left shift invariant, i.e. for any $f \in \mathcal{F}$ and $y \in X$ the function $x \mapsto y \cdot f(x)$ belongs to \mathcal{F} . Then $\mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I}) \setminus \mathcal{M}_\alpha(\mathcal{I}) \neq \emptyset$ for every $3 \leq \alpha < \omega_1$.*

Proof. Let $3 \leq \alpha < \omega_1$. Let A be a Π_α^0 set whose existence is guaranteed by Theorem 2.2.

Let h be a Borel function witnessing that \mathcal{F} is not \mathcal{I} -ubiquitous, i.e. there is no an \mathcal{I} -positive Borel set on which h equals to a function from \mathcal{F} .

Use Lemma 3.2 for A and h to find a $\Sigma_{\alpha+1}^0$ -measurable uniformization $f: \pi_1[A] \rightarrow X$ of A and a countable collection \mathcal{B} of pairwise disjoint Borel sets satisfying conditions (i)–(iii) of Lemma 3.2.

Suppose that $\text{graph}(f) \notin \mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I})$. Then, by the definition of $\mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I})$, for every $C \in \Sigma_{\alpha+2}^0$ with $C \supseteq \text{graph}(f)$ there is $g \in \mathcal{F}$ such that $\{x: g(x) \in C_x\} \notin \mathcal{I}$. Since $\text{graph}(f) \in \Sigma_{\alpha+2}^0$, there is $g \in \mathcal{F}$ with

$$\{x: g(x) \in \text{graph}(f)_x\} = \{x: g(x) = f(x)\} \notin \mathcal{I}.$$

Let $B' = \{x: g(x) = f(x)\}$. There is $B \in \mathcal{B}$ with $B' \cap B \notin \mathcal{I}$. Since $f|B = (y_B \cdot h)|B$, we have $g|(B' \cap B) = (y_B \cdot h)|(B' \cap B)$, and therefore $(y_B^{-1} \cdot g)|(B' \cap B) = h|(B' \cap B)$. As \mathcal{F} is left shift invariant, $y_B^{-1} \cdot g \in \mathcal{F}$, but this contradicts our assumption on h . Hence, $\text{graph}(f) \in \mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I})$.

By the choice of A for every $M \in \mathcal{M}_\alpha(\mathcal{I})$ there is $x \in \pi_1[A]$ such that $f(x) \notin M_x$. So, the graph of f is in $\mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I}) \setminus \mathcal{M}_\alpha(\mathcal{I})$. ■

Since $\mathcal{M}_\alpha(\mathcal{F}, \mathcal{I}) \subseteq \mathcal{M}_\alpha(\mathcal{I})$ for every $3 \leq \alpha < \omega_1$, the following corollary holds.

Corollary 3.4 *Consider \mathcal{I} and \mathcal{F} as in the above theorem. Then for every $3 \leq \alpha < \omega_1$ we have $\mathcal{M}_\alpha(\mathcal{F}, \mathcal{I}) \subsetneq \mathcal{M}_{\alpha+2}(\mathcal{F}, \mathcal{I})$. In particular, $\mathcal{M}_\alpha(\mathcal{I}) \subsetneq \mathcal{M}_{\alpha+2}(\mathcal{I})$.*

Corollary 3.5 *If \mathcal{J} is one of the following ideals of subsets of $[0, 1]^2$, then $\mathcal{J}_\alpha \subsetneq \mathcal{J}_{\alpha+2}$ for every $3 \leq \alpha < \omega_1$:*

- (1) *the ideal of sets A such that A can be covered by a Borel set whose all vertical sections are null (meager);*
- (2) *the ideal of sets A such that A can be covered by a Borel set whose all vertical and horizontal sections are null (meager);*
- (3) *the ideal of sets A such that A can be covered by a Borel set which is null (meager) in every direction;*
- (4) *the ideal of sets A such that A can be covered by a Borel set which is null (meager) on vertical sections and on graphs of polynomials.*

Proof. Consider $X = [0, 1)$ with addition modulo 1. Put $\mathcal{F}_{(1)} = \emptyset$, let $\mathcal{F}_{(2)}$ be the family of constant functions, $\mathcal{F}_{(3)}$ - the family of linear mappings and $\mathcal{F}_{(4)}$ - the family of polynomials. Then $\mathcal{J} = \mathcal{M}(\mathcal{F}_{(i)}, \mathcal{N}ull)$ (or $\mathcal{J} = \mathcal{M}(\mathcal{F}_{(i)}, \mathcal{M}eager)$). The result follows from Corollary 3.4. ■

Note that in fact, one can prove that $\mathcal{M}_\alpha(\mathcal{J}) \subsetneq \mathcal{M}_{\alpha+2}(\mathcal{J})$ assuming only that \mathcal{J} is a σ -ideal with a Borel base, containing all singletons, and which is Σ_α^0 -on- Π_α^0 and $\Sigma_{\alpha+1}^0$ -on- $\Pi_{\alpha+1}^0$. To see this, use Theorem 2.1 to find a $\Sigma_{\alpha+1}^0$ -measurable uniformization f of Π_α^0 set A from Theorem 2.2, and note that $\text{graph}(f) \in \mathcal{M}_{\alpha+2}(\mathcal{J}) \setminus \mathcal{M}_\alpha(\mathcal{J})$.

4 Fubini products

Notice that $\mathcal{M}(\mathcal{I})$ can be seen as $\{\emptyset\} \otimes \mathcal{I}$. Fubini products of null or meager ideals have bases of bounded Borel complexity (e.g. $\mathcal{N}ull \otimes \mathcal{N}ull$, $\mathcal{N}ull \otimes \mathcal{M}eager$, see e.g. [2]), so we cannot replace $\{\emptyset\}$ by $\mathcal{N}ull$ or $\mathcal{M}eager$ if we want to obtain a σ -ideal with the complex Borel base property. However, we will show that a σ -ideal of the form $\mathcal{J} \otimes \mathcal{N}ull$ ($\mathcal{J} \otimes \mathcal{M}eager$) has the complex Borel base property if \mathcal{J} has property (M) and a Borel base.

Definition 4.1 *We will say that an ideal \mathcal{J} of subsets of a Polish space X has property (M) if there is a Borel function $f: X \rightarrow [0, 1]$ such that $f^{-1}[\{x\}] \notin \mathcal{J}$ for every $x \in [0, 1]$.*

Notice that every uncountable Polish space X can play the role of $[0, 1]$ in the above theorem, since such X is Borel isomorphic to $[0, 1]$.

Let X be an uncountable Polish space and let $C \subseteq X$ be a set homeomorphic with the Cantor space $\{0, 1\}^\omega$. Since $\{0, 1\}^\omega$ is homeomorphic with $\{0, 1\}^\omega \times \{0, 1\}^\omega$, there is a continuous bijection $g: C \rightarrow \{0, 1\}^\omega \times \{0, 1\}^\omega$ and for every $x \in \{0, 1\}^\omega$ the preimage $g^{-1}[\{0, 1\}^\omega \times \{x\}]$ is uncountable. Then $f = \pi_2 \circ g$ witnesses that σ -ideal of countable subsets has property (M). Using a similar argument and the Silver Theorem [11, 35.20] one can show that ideals generated by an coanalytic equivalence relation with uncountable many equivalence classes (i.e. ideals of sets which can be covered by countably many equivalence classes) have property (M). By [7] the same holds for the ideal of subsets of the plane which can be covered by countably many lines. There are other σ -ideals with property (M): ideals on Polish spaces with a Σ_2^0 base which are not ccc (see [13]), some ideals defined by translations (see [3] and [14]). For the further discussion on property (M) see [1], [3] and [4].

Theorem 4.2 *Let \mathcal{I} be a σ -ideal of subsets of an uncountable Polish space X . Suppose \mathcal{I} has a Borel base, is Σ_α^0 -on- Π_α^0 for each $\alpha < \omega_1$ and contains all singletons. If a σ -ideal \mathcal{J} of subsets of X has property (M) then there is $\beta < \omega_1$ such that $(\mathcal{J} \otimes \mathcal{I})_\alpha \subsetneq (\mathcal{J} \otimes \mathcal{I})_{\alpha+2}$ for each $\alpha > \beta$.*

Proof. Let $f: X \rightarrow X$ be a Borel function witnessing (M) for \mathcal{J} , i.e. such that

$$f^{-1}[\{x\}] \notin \mathcal{J} \text{ for every } x \in X.$$

Let γ be such that f is Σ_γ^0 -measurable. Fix $\gamma < \beta < \omega_1$ such that $\gamma + \beta = \beta$ (e.g. $\beta = \omega^\gamma$) and notice that if $\alpha > \beta$ then $\gamma + \alpha = \alpha$. Let $\beta \leq \alpha < \omega_1$ and let $U \subseteq X \times X^2$ be universal for $\Sigma_\alpha^0(X^2)$. Define $H: X^3 \rightarrow X^3$ by $H(x, y, z) = (f(x), y, z)$ and let $V = H^{-1}[U]$. Clearly $V \in \Sigma_{\gamma+\alpha}^0 = \Sigma_\alpha^0$. Consider $A = \{(x, y) \in X^2: (x, x, y) \notin V\}$, a Π_α^0 set.

Since \mathcal{I} is Σ_α^0 -on- Π_α^0 , the set $A' = A \cap (\{x: A_x^c \in \mathcal{I}\} \times X)$ is also Π_α^0 . By Theorem 2.1, the set A' has a uniformization with a graph C of class $\Pi_{\alpha+1}^0$. Clearly $C \in \mathcal{M}_{\alpha+2}(\mathcal{I})$.

We will show that $C \notin (\mathcal{J} \otimes \mathcal{I})_\alpha$ by proving that $C \not\subseteq E$ for $E \in \Sigma_\alpha^0(X^2) \cap (\mathcal{J} \otimes \mathcal{I})$. Of course $E = U_x$ for some $x \in X$. So, $E = V_t$ for each $t \in f^{-1}[\{x\}]$. Notice that $C_t \cap E_t = \emptyset$ since $E_t = (V_t)_t = V_{(t,t)} = [0, 1] \setminus A_t$. By the definition of Fubini product $\{t \in f^{-1}[\{x\}]: E_t \notin \mathcal{I}\} \in \mathcal{J}$. As $f^{-1}[\{x\}] \notin \mathcal{J}$, we have also that $\{t \in f^{-1}[\{x\}]: E_t \in$

$\mathcal{I}\} \notin \mathcal{J}$. Moreover, if $t \in f^{-1}[\{x\}]$ and $E_t \in \mathcal{I}$ then $A_t^c = E_t \in \mathcal{I}$. Hence there is t such that $(A_t) \cap E_t = \emptyset$ and $A_t \supseteq C_t \neq \emptyset$. Therefore $C \setminus E \neq \emptyset$, which means that $C \notin (\mathcal{J} \otimes \mathcal{I})_\alpha$. Since $\mathcal{M}_{\alpha+2}(\mathcal{I}) \subseteq (\mathcal{J} \otimes \mathcal{I})_{\alpha+2}$, it follows that $C \in (\mathcal{J} \otimes \mathcal{I})_{\alpha+2} \setminus (\mathcal{J} \otimes \mathcal{I})_\alpha$. ■

In [6], the σ -ideals $\mathcal{M}(\mathcal{N}ull)$ and $\mathcal{M}(\mathcal{M}eager)$ are called Mokobodzki σ -ideals. The results presented here and in the previous section indicate how we can generalize the definition of a Mokobodzki ideal.

Let (X, \cdot) be an uncountable Polish group. We say that $Y \subseteq X^2$ belongs to a σ -ideal $\mathcal{M}(\mathcal{F}, \mathcal{I}, \mathcal{J})$ on X^2 if $Y \in \mathcal{J} \otimes \mathcal{I}$ and Y can be covered by a Borel set $B \subseteq X^2$ such that

$$\{y \in X : \{x \in X : (x, y \cdot f(x)) \in B\} \notin \mathcal{I}\} \in \mathcal{J}$$

for every $f \in \mathcal{F}$. We say that a σ -ideal is a *generalized Mokobodzki ideal* if it is of the form $\mathcal{M}(\mathcal{F}, \mathcal{I}, \mathcal{J})$ for $\mathcal{I} = \mathcal{N}ull$ or $\mathcal{I} = \mathcal{M}eager$, a family of Borel functions $\mathcal{F} \subseteq X^X$ containing constants which is left shift invariant and is not \mathcal{I} -ubiquitous, and a σ -ideal \mathcal{J} with property (M). Loosely speaking, a generalized Mokobodzki ideal is a σ -ideal consisting of sets C which are null (meager) along all functions from \mathcal{F} and which have null (meager) vertical section C_x for \mathcal{J} -almost every x . By combining proofs of Theorem 3.3 and Theorem 4.2 it is possible to show that all generalized Mokobodzki ideals have the complex Borel base property. We decided to prove it in less general cases for the sake of clarity.

5 Set-theoretic properties

In this section we explore some properties of the ideals considered in the previous sections. First, we show that they have property (M). As before, throughout this section we assume that X is an uncountable Polish space.

Proposition 5.1 *If \mathcal{I} is a σ -ideal of subsets of X such that $X \notin \mathcal{I}$ and $\mathcal{F} \subseteq X^X$, then $\mathcal{M}(\mathcal{F}, \mathcal{I})$ has property (M). Also, if a σ -ideal \mathcal{J} has property (M), then $\mathcal{J} \otimes \mathcal{I}$ has property (M).*

Proof. Consider the function $\pi_1(x, y) = x$. Then $\pi_1^{-1}[\{x\}] = \{x\} \times X \notin \mathcal{M}(\mathcal{F}, \mathcal{I})$. To prove the second part of the theorem consider $g = f \circ \pi_1$, where $f: X \rightarrow X$ witnesses that \mathcal{J} has property (M). ■

Notice, that this means that the ideals from the previous sections are very far from being ccc. Notice also that the above proposition together with Theorem 4.2 allows us to produce easily a lot of examples of σ -ideals with the complex Borel base property. In particular, Fubini products of $\mathcal{N}ull$ or $\mathcal{M}eager$ ideals with ideals from Proposition 5.1 have the complex Borel base property.

In [6, 2.4] it was proved that if \mathcal{I} a proper σ -ideal of subsets of X , then $\mathcal{M}(\mathcal{I})$ has additivity ω_1 . The proof uses Theorem 2.2, Kondo–Adisson and Sierpiński theorems.

In fact, it seems that the essential reason for the equality $\text{add}(\mathcal{M}(\mathcal{I})) = \omega_1$ lies in the fact that $\mathcal{M}(\mathcal{I})$ is an union of sequences of ω_1 strictly increasing σ -ideals, $\mathcal{I} \in \{\mathcal{N}ull, \mathcal{M}eager\}$. More generally, the following fact holds.

Proposition 5.2 *Suppose that \mathcal{I} has a Borel base and there are σ -ideals \mathcal{I}_α with $\mathcal{I} = \bigcup_{\alpha < \kappa} \mathcal{I}_\alpha$ and $\mathcal{I}_\alpha \subsetneq \mathcal{I}_{\alpha+1}$ for every $\alpha < \kappa$. Then $\text{add}(\mathcal{I}) \leq \kappa$. In particular, if \mathcal{I} has the complex Borel base property, then $\text{add}(\mathcal{I}) = \omega_1$.*

Proof. For every $\alpha < \kappa$ take $A_\alpha \in \mathcal{I}_{\alpha+1} \setminus \mathcal{I}_\alpha$ and consider a set $A = \bigcup_{\alpha < \omega_1} A_\alpha$. If A is in \mathcal{I} , then we could find Borel $B \in \mathcal{I}$ with $B \supseteq A$, but there would be $\alpha < \kappa$ with $B \in \mathcal{I}_\alpha$. This would contradict the fact that $A_\alpha \subseteq B$ and $A_\alpha \notin \mathcal{I}_\alpha$.

The second part of the statement follows easily. ■

As a corollary we obtain that the ideals from Theorem 4.2 and Theorem 3.3 (and, all generalized Mokobodzki ideals) have additivity ω_1 . In fact, every σ -ideal containing all singletons is a union of a strictly increasing sequence of its proper sub- σ -ideals.

Lemma 5.3 *If \mathcal{I}, \mathcal{J} are σ -ideals on X and $[X]^\omega \subseteq \mathcal{J} \subsetneq \mathcal{I}$, then there is $A \in \mathcal{I} \setminus \mathcal{J}$ such that the σ -ideal generated by \mathcal{J} and A is not equal to \mathcal{I} .*

Proof. First, notice that for every $\varepsilon > 0$ and every $A \in \mathcal{I} \setminus \mathcal{J}$ there is $B \subseteq A$ of diameter at most ε such that $B \in \mathcal{I} \setminus \mathcal{J}$. It follows from the fact that A is a countable union of sets with diameter at most ε .

Now, construct a decreasing sequence B_1, B_2, \dots such that $B_n \in \mathcal{I} \setminus \mathcal{J}$ and $\text{diam}(B_n) < 1/n$. Let $A_n = B_n \setminus B_{n+1}$. If $A_n \in \mathcal{J}$ for each n , then $\bigcup A_n \in \mathcal{J}$. Since $\bigcup A_n = B_1 \setminus \bigcap B_n$ and $|\bigcap B_n| \leq 1$, then $B_1 \in \mathcal{J}$, a contradiction. So, there is n such that $A_n \in \mathcal{I} \setminus \mathcal{J}$ and $B_{n+1} \in \mathcal{I} \setminus \mathcal{J}$. The σ -ideal generated by \mathcal{J} and A_n does not contain B_{n+1} and therefore is not equal to \mathcal{I} . ■

Using the above fact for every σ -ideal \mathcal{I} strictly extending $[X]^\omega$ one can easily construct a strictly increasing sequence of σ -ideals $(\mathcal{I}_\alpha)_{\alpha < \beta}$ such that $\mathcal{I}_\alpha \neq \mathcal{I}$ for each α (in fact, one can easily construct such a sequence also for $[X]^\omega$, enumerating X and considering σ -ideals generated by the initial segments). So, we can define a coefficient indicating what is the minimal size of such a sequence: $\text{cofin}(\mathcal{I}) = \min\{\kappa: \mathcal{I} = \bigcup_{\alpha < \kappa} \mathcal{I}_\alpha \text{ where } (\mathcal{I}_\alpha) \text{ is a strictly increasing sequence of } \sigma\text{-ideals which are proper subideals of } \mathcal{I}\}$. Clearly $\text{cofin}(\mathcal{I})$ is regular. Moreover, an argument as in the proof of Proposition 5.2 implies that $\text{add}(\mathcal{I}) \leq \text{cofin}(\mathcal{I})$. If \mathcal{I} has the complex base property then $\text{cofin}(\mathcal{I}) = \omega_1$. It is interesting how this coefficient behaves in other situations. In the last section we pose one of the natural questions in this context.

We will turn now to classical ideal invariants. Some cardinal coefficients of $\mathcal{M}(\mathcal{F}, \mathcal{I})$ are inherited from the ideal \mathcal{I} .

Proposition 5.4 *If \mathcal{I} is a σ -ideal of subsets of X and $\mathcal{F} \subseteq X^X$, then*

(i) $\text{cov}(\mathcal{M}(\mathcal{F}, \mathcal{I})) = \text{cov}(\mathcal{I})$ provided \mathcal{I} has a Borel base;

(ii) $\text{non}(\mathcal{M}(\mathcal{F}, \mathcal{I})) = \text{non}(\mathcal{I})$.

Proof. Suppose that the family $\{A_\xi: \xi < \kappa\} \subseteq \mathcal{I}$ covers X . We may assume that each A_ξ is Borel. Then $\{A_\xi \times A_\eta: \xi, \eta < \kappa\} \subseteq \mathcal{M}(\mathcal{F}, \mathcal{I})$ and it covers X^2 . On the other hand if $\{Z_\xi: \xi < \kappa\} \subseteq \mathcal{M}(\mathcal{F}, \mathcal{I})$ covers X^2 and $x \in X$, then $\{(Z_\xi)_x: \xi < \kappa\} \subseteq \mathcal{I}$ and it covers X .

If $Z \notin \mathcal{I}$, then $\{x\} \times Z \notin \mathcal{M}(\mathcal{F}, \mathcal{I})$ and $|\{x\} \times Z| = |Z|$, where $x \in X$. On the other hand, if $|Z| < \text{non}(\mathcal{I})$ for $Z \subseteq X^2$, then $|\pi_1[Z]| < \text{non}(\mathcal{I})$ and $|\pi_2[Z]| < \text{non}(\mathcal{I})$. So, there is a Borel $B \in \mathcal{I}$ such that $\pi_1[Z] \cup \pi_2[Z] \subseteq B$. Therefore, $Z \subseteq B \times B$ and $Z \in \mathcal{M}(\mathcal{F}, \mathcal{I})$. ■

The assumption that \mathcal{I} has a Borel base in part (i) of the assertion of Proposition 5.4 is needed. To see this, let $\{B_\xi: \xi < \omega_1\}$ be a collection of pairwise disjoint Bernstein sets in $[0, 1]$ which covers $[0, 1]$. Define \mathcal{I} as the σ -ideal of subsets of $[0, 1]$ which can be covered by countably many sets B_ξ . Then $\text{cov}(\mathcal{I}) = \omega_1$. But $\mathcal{M}(\mathcal{I})$ consists of sets with countable sections, hence $\text{cov}(\mathcal{M}(\mathcal{I})) = \mathfrak{c}$ and consistently $\text{cov}(\mathcal{M}(\mathcal{I})) > \text{cov}(\mathcal{I})$.

In the case of ideals defined in Section 4 we can apply the following fact.

Proposition 5.5 (folklore) *If \mathcal{I} and \mathcal{J} are σ -ideals of subsets of X with Borel bases then*

$$\text{non}(\mathcal{J} \otimes \mathcal{I}) = \max\{\text{non}(\mathcal{I}), \text{non}(\mathcal{J})\} \text{ and}$$

$$\text{cov}(\mathcal{J} \otimes \mathcal{I}) = \min\{\text{cov}(\mathcal{I}), \text{cov}(\mathcal{J})\}.$$

Proof. Let Z, Y witness $\text{non}(\mathcal{J})$ and $\text{non}(\mathcal{I})$ respectively. Then $Z \times Y \notin \mathcal{J} \otimes \mathcal{I}$ and $|Z \times Y| = \max\{\text{non}(\mathcal{I}), \text{non}(\mathcal{J})\}$. If Z witnesses $\text{non}(\mathcal{J} \otimes \mathcal{I})$ then $|\pi_1[Z]| \geq \text{non}(\mathcal{J})$ and $|\pi_2[Z]| \geq \text{non}(\mathcal{I})$. Therefore, $\text{non}(\mathcal{J} \otimes \mathcal{I}) = \max\{\text{non}(\mathcal{I}), \text{non}(\mathcal{J})\}$.

Assume that $\{A_\xi: \xi < \kappa\} \subseteq \mathcal{J}$ covers X . Then, $\{A_\xi \times X: \xi < \kappa\} \subseteq \mathcal{J} \otimes \mathcal{I}$ and it covers X^2 . A similar argument for \mathcal{I} shows that $\text{cov}(\mathcal{J} \otimes \mathcal{I}) \leq \min\{\text{cov}(\mathcal{I}), \text{cov}(\mathcal{J})\}$. On the other hand, if $\text{cov}(\mathcal{J} \otimes \mathcal{I}) \leq \text{cov}(\mathcal{J})$ and $\{Z_\xi: \xi < \text{cov}(\mathcal{J} \otimes \mathcal{I})\}$ covers X^2 , then there is $x \in X$ such that $(Z_\xi)_x \in \mathcal{I}$ for every $\xi < \kappa$. As $(Z_\xi)_x$ covers X , we have $\text{cov}(\mathcal{J} \otimes \mathcal{I}) \geq \min\{\text{cov}(\mathcal{I}), \text{cov}(\mathcal{J})\}$. ■

Let \mathcal{I} be a σ -ideal on a Polish space X . Recall that $H \subseteq X$ is an \mathcal{I} -hull of a set $A \subseteq X$ if H is Borel and for every Borel $B \subseteq H$ either $B \in \mathcal{I}$ or $B \cap A \neq \emptyset$. We say that a σ -ideal \mathcal{I} on X has the hull property if every subset of X has an \mathcal{I} -hull.

Proposition 5.6 *If \mathcal{I} is a Borel-on-Borel σ -ideal on X , then $\mathcal{M}(\mathcal{I})$ does not possess the hull property.*

Proof. Consider $A = V \times X$, where $V \subseteq X$ is a Bernstein set. Assume $H \supseteq A$ is a Borel set and consider $L = \{x \in X: H_x \notin \mathcal{I}\}$. Notice that L is Borel. Since $V \subseteq L$, the set L has to be co-countable. In particular, there is $x \in L \setminus V$ and the set $\{x\} \times H_x$ is an $\mathcal{M}(\mathcal{I})$ -positive Borel subset of H disjoint with A . ■

6 Additional remarks and open questions

It is well known that every Borel function on $[0, 1]^2$ is nice on a big domain (of Baire class 1 on a set of Lebesgue measure 1, continuous on a co-meager set). However, in [2] it is proved that this is not the case if instead of $\mathcal{N}ull$ or $\mathcal{M}eager$ ideals we would consider $\mathcal{M}(\mathcal{N}ull)$ or $\mathcal{M}(\mathcal{M}eager)$. More precisely, for every $\alpha \geq 2$ there is a Borel function $g: [0, 1]^2 \rightarrow [0, 1]$ such that for every Borel set $M \in \mathcal{M}(\mathcal{N}ull)$ (or $\mathcal{M}(\mathcal{M}eager)$) we can find $x \in [0, 1]$ such that the function $g_x|_{M_x^c}$ is not Σ_α^0 -measurable (here $g_x(y) = g(x, y)$). On the other hand (see [2]), these pathological points can be covered by null and meager sets, i.e. for every M as above, the set

$$\{x: g_x|_{M_x^c} \text{ is not } \Sigma_\alpha^0\text{-measurable}\} \in \mathcal{N}ull \cap \mathcal{M}eager.$$

The following notion is motivated by the above considerations.

Let X be an uncountable Polish space. Let \mathcal{J}, \mathcal{I} be σ -ideals of subsets of X . We say that \mathcal{J} is \mathcal{I} -thin if for each $2 \leq \alpha < \omega_1$ there is a Borel function $g: X^2 \rightarrow X$ such that for all $M \in \mathcal{M}_\alpha(\mathcal{I})$ and $C \in \mathcal{J}$ there is $t_0 \in X \setminus C$ such that $g_{t_0}|_{(X \setminus M_{t_0})}$ is not Σ_α^0 -measurable. Using the notion of thinness, the theorems cited above can be stated in this form: $\{\emptyset\}$ is $\mathcal{N}ull$ -thin and $\mathcal{M}eager$ -thin, whereas $\mathcal{N}ull$ and $\mathcal{M}eager$ are not. We show the following.

Proposition 6.1 *Every σ -ideal with property (M) is \mathcal{I} -thin for every σ -ideal \mathcal{I} that is Borel-on-Borel.*

Proof. Let \mathcal{I} be a Borel-on-Borel σ -ideal on X and let \mathcal{J} be a σ -ideal on X with the property (M). Assume that $2 \leq \alpha < \omega_1$. Let $f: X \rightarrow X$ be a Borel function witnessing (M) for \mathcal{J} . Let $U \subseteq X \times X^2$ be universal for $\Sigma_\alpha^0(X^2)$. As in the proof of Theorem 4.2 define $H: X^3 \rightarrow X^3$ by $H(x, y, z) = (f(x), y, z)$, let $V = H^{-1}[U]$ and consider $A = \{(x, y) \in X^2: (x, x, y) \notin V\}$. Notice that A is Borel, and thus the set

$$A' = A \cap \{(t, y): A_t^c \in \mathcal{I}\}$$

is Σ_γ^0 for some $\gamma < \omega_1$.

Let $F: \{0, 1\}^\omega \times X \rightarrow X$ be a function from Theorem 2.1 for the set A' and the ordinal γ . Define $G: \{0, 1\}^\omega \times X \rightarrow X^2$ by

$$G(c, x) = (x, F(c, x)).$$

Consider a Borel subset D of $\{0, 1\}^\omega$ such that $D \notin \Sigma_\alpha^0(\{0, 1\}^\omega)$ and let

$$\tilde{D} = G[D \times \pi_1[A']].$$

Notice that since G is one-to-one and Borel, the set \tilde{D} is Borel. Clearly $\tilde{D} \subseteq A'$. Let $g: X^2 \rightarrow \{x_0, x_1\}$ be the characteristic function of \tilde{D} , where x_0 and x_1 are distinct points of X .

Let $M \in \mathcal{M}_\alpha(\mathcal{I}) \cap \Sigma_\alpha^0$ and $C \in \mathcal{J}$. There is $x \in X$ such that $U_x = M$. So, for every $t \in f^{-1}[\{x\}]$ we have $A_t = M_t^c$ and, therefore, $t \in \pi_1[A']$. Since $f^{-1}[\{x\}] \setminus C \neq \emptyset$, there is t_0 with

$$(g_{t_0}|_{(X \setminus M_{t_0})})^{-1}[\{x_1\}] = (X \setminus M_{t_0}) \cap \tilde{D}_{t_0} = \tilde{D}_{t_0} = G[D \times \{t_0\}].$$

Since $c \mapsto G(c, t_0)$ is continuous and one-to-one, the set $G[D \times \{t_0\}]$ is homeomorphic to D . Hence $g_{t_0}|_{(X \setminus M_{t_0})}$ is not Σ_α^0 -measurable.

■

We finish with a list of open questions.

Problem 6.2 *Suppose X is an uncountable Polish space and \mathcal{I} is a σ -ideal of subsets of X .*

- (i) *Is it true that $\mathcal{M}_\alpha(\mathcal{I}) \subsetneq \mathcal{M}_{\alpha+1}(\mathcal{I})$ or even $\mathcal{M}_\alpha(\mathcal{F}, \mathcal{I}) \subsetneq \mathcal{M}_{\alpha+1}(\mathcal{F}, \mathcal{I})$ for every $\alpha < \omega_1$, $\mathcal{I} \in \{\mathcal{N}ull, \mathcal{M}eager\}$ and every family \mathcal{F} as in Theorem 3.3?*
- (ii) *Is it provable in ZFC that for every \mathcal{I} containing all singletons we have $\text{add}(\mathcal{I}) = \text{cofin}(\mathcal{I})$?*
- (iii) *Does there exist \mathcal{I} with the hull property and the complex Borel base property?*
- (iv) *Let $\mathcal{K}([0, 1]^2)$ stands for the set of all nonempty compact subsets of $[0, 1]^2$ with the Vietoris topology. What is the complexity of the set $\{K \in \mathcal{K}([0, 1]^2) : K \in \mathcal{M}(\mathcal{I})\}$ for $\mathcal{I} \in \{\mathcal{N}ull, \mathcal{M}eager\}$? Is it Π_1^1 -complete?*

References

- [1] M. Balcerzak, *Can ideals without ccc be interesting?* Topology Appl. **55** (1994), no. 3, 251–260;
- [2] M. Balcerzak and Sz. Głab, *Measure-category properties of Borel plane sets and Borel functions of two variables*, Acta Math. Hungar., **126** (3) (2010), 241–252;
- [3] M. Balcerzak, A. Rosłanowski and S. Shelah, *Ideals without ccc*, J Symbolic Logic **63** (1998), 128–147;
- [4] H. Becker, *Ideals without ccc and without property (M)*, Proc. Amer. Math. Soc. **128** (2000), no. 10, 3031–3034;
- [5] A. Blass, *Combinatorial cardinal characteristics of the continuum*, to appear as a chapter in the *Handbook of Set Theory*;
- [6] J. Cichoń and J. Pawlikowski, *On ideals of subsets of the plane and on Cohen reals*, The Journal of Symbolic Logic **51** (1986), no. 3, 561–569;
- [7] F. van Engelen, K. Kunen, A. W. Miller, *Two remarks about analytic sets*, Set theory and its applications (Toronto, ON, 1987), 68–72, Lecture Notes in Math., 1401, Springer, Berlin, 1989;
- [8] I. Farah and J. Zapletal, *Between Maharam and von Neumann’s problems*, Math. Reseach Letters **11** (2004), no 5–6, 673–684;
- [9] D. Fremlin, *Consequences of Martin’s Axiom*, Cambridge Tracts in Math. 84, Cambridge University Press (1984);
- [10] P. Holický, *Borel classes of uniformizations of sets with large sections*, Fund. Math. **207** (2010), 145–160;

- [11] A. Kechris, *Classical descriptive set theory*, Springer-Verlag, New York, 1995;
- [12] K. Kuratowski, *Topology* Vol. 1, Academic Press, New York and London, 1966;
- [13] I. Reclaw, P. Zakrzewski, *Fubini properties of ideals*, Real Anal. Exchange **25** (1999/00), no. 2, 565–578;
- [14] S. Solecki, *A Fubini theorem*, Topology Appl. **154** (2007), no. 12, 2462–2464;
- [15] P. Zakrzewski, *On the uniqueness of measure and category σ -ideals on 2^ω* , J. Appl. Anal. **13(2)** (2007), 249–257.

Instytut Matematyczny, Uniwersytet Wrocławski
plac Grunwaldzki 2/4, 50-384 Wrocław
PBOROD@MATH.UNI.WROC.PL
<http://www.math.uni.wroc.pl/~pborod>

Institute of Mathematics, Technical University of Łódź
ul. Wólczańska 215, 90-924 Łódź
SZYMON2377@O2.PL
<http://im0.p.lodz.pl/~sglab/indexeng.html>